

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
12 April 2001 (12.04.2001)


PCT

(10) International Publication Number
WO 01/25454 A2

- (51) International Patent Classification⁷: C12N 15/82, 9/02 (74) Agents: CÔTÉ, France et al.; Swabey Ogilvy Renault, Suite 1600, 1981 McGill College Avenue, Montréal, Québec H3A 2Y3 (CA).
- (21) International Application Number: PCT/CA00/01143
- (22) International Filing Date: 2 October 2000 (02.10.2000) (81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data: 60/157,133 4 October 1999 (04.10.1999) US
- (71) Applicant (*for all designated States except US*): MED-ICAGO INC. [CA/CA]; 2480, rue Hochelaga, Sainte-Foy, Québec G1K 7P4 (CA).
- (72) Inventors; and
- (75) Inventors/Applicants (*for US only*): VÉZINA, Louis-Philippe [CA/CA]; 206, Route 138, Neuville, Québec G0A 2R0 (CA). D'AOUST, Marc-André [CA/CA]; 939, avenue Manrèse, App. 2, Québec, Québec G1S 2W9 (CA).
- (84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).
- Published:
— Without international search report and to be republished upon receipt of that report.

[Continued on next page]

(54) Title: METHOD FOR REGULATING TRANSCRIPTION OF FOREIGN GENES



(57) Abstract: The present invention relates to a method of regulating the transcription of transgene in genetically-modified organisms. More specifically, the invention relates to the use of expression vectors harboring the coding sequence of a gene of interest under the transcriptional control of promoting sequences for which activity is regulated by the presence of nitrogen.

WO 01/25454 A2



For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

METHOD FOR REGULATING TRANSCRIPTION OF FOREIGN GENES

BACKGROUND OF THE INVENTION

(a) Field of the Invention

5 The invention relates to a method of regulating the transcription of transgene in genetically-modified organisms. More specifically, the invention relates to the use of expression vectors harboring the coding sequence of a gene of interest under the transcriptional control of promoting sequences for which activity is regulated by the presence of
10 nitrogen. Preferably, these constructs are used in transgenic leguminous plants (for example soybean, alfalfa, clover, birdsfoot trefoil, beans, peas, peanuts) where growth is not impaired by lack of mineral nitrogen, and in which induction of expression could be performed at any given time during development, through the addition of a suitable nitrogen source. In a
15 broader perspective, the invention could be used to induce expression of any given transgene through the addition of any nitrogen source, provided that the organism can be grown adequately in the absence of this nitrogen inducer; as an example within the plant kingdom, duckweed (*Lemna minor*) can adapt to grow either on nitrate or ammonium as nitrogen
20 source; transgenic duckweed could therefore be grown on nitrate as a sole nitrogen source and expression of the transgene triggered by the addition of ammonium, provided that the cassette contains a promoter from a native gene for which expression is turned on by the addition of ammonium. The invention therefore provides a means of regulating the
25 expression of a transgenic trait in any organism through the addition of various nitrogenous inducer.

(b) Description of Prior Art

 Nitrogen is a molecule essential to life. All living organism need nitrogen in order to synthesize amino acids, the building blocks of
30 proteins, and nucleotides, the building blocks of nucleic acids. It is Ammonium nitrate is the preferred form of mineral nitrogen provided to crops in the form of fertilizer. Nitrate-nitrogen is first reduced to nitrite and then to ammonium through the activity of a metabolic pathway common to most herbaceous plants. Depending on the species, part or all of the
35 absorbed nitrate will move to leaf cells through the xylem before it is

reduced to ammonium. Ammonium, or other reduced forms of nitrogen are also absorbed (although usually at lower rates) by the root system but their assimilation does not require reduction. These newly absorbed ammonium or ammonium-containing molecules join the endogenous
5 pools in the cells which is formed by ammonium cycling through amino acids and other nitrogenous molecules. Some species do not metabolize nitrate-nitrogen easily and therefore cannot rely on nitrate as sole nitrogen source; many coniferous species fall into this latter category. Legumes and other symbiotic plant species form a third large class of nitrogen user
10 within the plant kingdom; they form a metabolic alliance with a microbial organism through which they can fix gaseous nitrogen. This reduced nitrogen is used efficiently by the plant for growth, and therefore, these crops can develop independently of the availability of mineral nitrogen in the soil.

15 Many microbes and wild plant species will adapt extensively to availability of nitrogen sources and can therefore complete their life cycle in the absence of one molecular form of nitrogen, which they could use exclusively and efficiently if available in another growing environment. As for most assimilatory pathways, nitrogen assimilation is tightly regulated in
20 cells. As an example, the expression of genes encoding nitrate reductase (NaR) and nitrite reductase (NiR), which are responsible for the reduction of nitrate to ammonium, has been extensively described in various microbial and plant species (for a review, see Mifflin and Lea, Books 5 and 12, in *The Biochemistry of plants*). Although nitrate is not the only
25 regulatory molecule involved in the control of NaR and NiR expression, its presence is essential to initiate the cascade of transduction that eventually leads to sustained transcription and translation of these genes. It has been shown that expression of NaR and NiR genes is repressed in leguminous plants when they are grown in the absence of mineral
30 nitrogen

NiR promoters have been characterized in some plant species (Back et al., 1991, *Plant Molecular Biology* 17:9-18; Sander et al., 1995, *Plant Molecular Biology* 27:165-177). Inducibility of these promoters have also been characterized using marker genes in transgenic plants, where it

was shown that availability of nitrate is required for full activation of transcription.

Assimilatory pathways for other nitrogen sources have also been described, and promoters for genes involved in some of these pathways have also been characterized.

Genetic transformation of microbes have been used for more than 15 years to produce useful recombinant molecules, and applications in the pharmaceutical, cosmaceutical and dermaceutical industries are being currently exploited. This technology has expanded from microbes to plants and animals in the last ten years with the development of techniques required to adapt this general concept to complex eukaryotic organisms. Basically a gene encoding for a protein of interest or a gene encoding for an enzyme responsible for a modification of a metabolic pathway that leads to a molecule of interest, is linked in an appropriate fashion to cis-and trans-acting regulatory sequences, and transferred to a target cell where it is incorporated in the molecular machinery (in a transitory or stable fashion). The transgenic cell, or a tissue or organism regenerated from the transgenic cell will then perform transcription and translation of the transgene and therefore be enabled to accumulate the protein of interest or to perform the new metabolic reaction through the activity of the enzyme of interest.

The emerging industry of molecular farming (production of recombinant molecules in animals or crops) is one of the most promising industry of the coming century. Its promise is to provide safe and renewable molecule factories for the industry. Among the applications that are currently developed are the production of low-cost monoclonal antibodies for therapeutic and diagnostic uses, the production of unlimited amounts of hormones, cytokines and other bio-active molecules for the treatment of chronicle or lethal diseases, the production of bio-safe substitutes for various blood components, the production of unlimited amounts of processing enzymes for the food and pulp industry, the production of low-cost enzymes for waste treatments, and the production of safe bio-active molecules for the cosmetic industry.

Limitations to the application of this technology has often come from the inability of transgenic organisms to accumulate adequate

amounts of the recombinant product, as a result of low transcription rates, improper splicing of the messenger, instability of the foreign mRNA, poor translation rates, hyper-susceptibility of the recombinant protein to the action of endogenous proteases or hyper-susceptibility of the recombinant organism to the foreign protein which result in improper and limited growth or in the worst cases, in strong deleterious effects to the host organism. Inadequacy of production level has a direct impact on the development of applications when profit margins are narrow, or when treatment and/or disposal of residual matter causes bio-safety or environmental problems. Improvement of the accumulation level of the desired recombinant product thus appears to be one critical factor that warrants commercialization of many applications of molecular farming.

The use of inducible promoters has been proposed, and in some instances used successfully, to counteract the combined effect of all the above-mentioned factors. Strong inducible promoters may succeed in generating high ephemeral transcription rates which result in high transitory accumulation of foreign mRNA and translational product. As a result, when inducibility of expression is paired with adequate synchronized protein recovery procedures, the yield per unit obtained is higher than with the use of constitutive expression.

Several expression cassettes harboring inducible promoters have been developed for microbial production systems, and some are currently available for research purposes. Some inducible promoters are currently used in plant (wound inducibility) or animal (specificity to cells of the mammary glands, PPL) systems, although none reported are using low-cost and bio-safe chemical inducers such as nitrate salts.

It would be highly desirable to be provided with a method of regulating the transcription of transgene in genetically-modified organisms.

30

SUMMARY OF THE INVENTION

One aim of the present invention is to provide a method of regulating the transcription of transgene in genetically-modified organisms.

Another aim of the present invention is to provide the use of expression vectors harboring the coding sequence of a gene of interest under the transcriptional control of promoting sequences for which activity is regulated by the presence of nitrogen.

5 The present invention relates to the use of a nitrogen-inducible expression cassettes for the controlled expression of foreign genes in plants. It will be shown from the following description that isolating such a regulatory sequences can be performed so that when cis-acting sequences are appropriately associated to the open reading frame of a
10 gene of interest, its transcription can be controlled by the addition of specific nitrogen sources.

 In one aspect of this invention, the targeted system uses leguminous plant species, so that constructs containing a nitrate-inducible promoter will be maintained transcriptionally low throughout the growth
15 period if the transgenic plant is maintained on a nitrate-free medium, thus allowing the development of the plant biomass without interference from the transgenic trait. Upon addition of nitrate to the growth medium, transcription will be induced in a relatively large proportion of the biomass over the following days. Optimization of induction time and protein
20 accumulation will then be performed in order to maximize recovery of the desired recombinant product.

 Although the following description will make clear that this invention can be easily adapted to nitrate induction on nitrate-deprived transgenic leguminous plants, it should be remembered that this general
25 concept can also be applied to the development of other production systems, making profit of the wide variety of nitrogen assimilation systems in the microbial, plant and animal kingdoms.

 In one other aspect of this invention, nitrogen inducibility can also be used to maximize protein production in organisms which do not
30 perform nitrogen fixation through symbiotic association, but that can use variable sources of nitrogen (reduced or oxidized) for growth, and thus possess the ability to develop adequately while one of their nitrogen assimilation pathway is inactive due to lack of one nitrogenous substrate in the growing environment. Using an expression cassettes that controls
35 the transcription of any gene in this inactive pathway in order to drive the

expression of a gene of interest in such an organism, will allow for inducible expression of the transgenic trait upon addition of the previously lacking nitrogenous compound. As an example, duckweed is a plant species that can grow alternately on nitrate or ammonium; this invention
5 could be used to develop an expression cassette harboring an ammonium-inducible promoter appropriately linked to a gene of interest so that the induction would be performed on nitrate-grown transgenic duckweed plants.

In accordance with the present invention there is provided a
10 method for regulating transcription of a foreign gene in transgenic organisms, comprising the steps of:

- a) preparing a transgenic organism using an expression construct consisting of at least a nitrogen-inducible promoter having a sequence selected from the group consisting of SEQ ID NOS: 1
15 to 13 and functional fragments and derivatives thereof, and an ORF of a gene, wherein said promoter is operationally located with respect to said gene for expression of said gene.

In accordance with a preferred embodiment of the present invention, the method of may further comprise the step of regulating
20 transcriptional expression of said gene by addition or removal of a nitrogen inducer.

In accordance with a preferred embodiment of the present invention, there is provided a method for regulating transcription of a foreign gene in transgenic organisms comprising:

- 25 a) preparing an expression construct consisting of at least a nitrogen-inducible promoter with or without cis-acting sequence, an ORF of a gene, and a polyadenylation signal end site at the 3'end of said construct, wherein said promoter is operationally located with respect to said gene for expression of said gene
30 and modulated for transcriptional expression of said gene by addition or removal of a nitrogen inducer;
- b) sub-cloning the construct of step a) into a suitable transfection vector for said organism;
- c) transferring said vector into DNA of said organism or a cell
35 thereof; and

- d) selecting for transgenicity on a suitable medium.

In accordance with a preferred embodiment of the present invention, the method of may further comprise the steps of:

- 5 e) introducing the vector into a suitable *Agrobacterium tumefaciens* strain;
- f) using the *Agrobacterium* strain of step a) to transfer T-DNA into a plant cell;
- 10 g) selecting for transgenicity of said plant cell on a suitable medium;
- h) regenerating embryos or plantlets from said transgenic cells; and
- i) growing mature plants from said regenerated embryos.

15 In accordance with a preferred embodiment of the present invention, the cis-acting sequence may be isolated from 5' upstream region of an expressed Nir gene in alfalfa.

In accordance with a preferred embodiment of the method of the present invention, the promoter has the sequence set forth in SEQ ID NO:1 to 13 and functional fragments and derivatives thereof.

20 In accordance with a preferred embodiment of the present invention, the organism is a plant, more preferably a dicotyledonous plant.

In accordance with a preferred embodiment of the present invention, the organism is alfalfa or tobacco.

25 In accordance with a preferred embodiment of the present invention, the nitrogen inducer is nitrate.

In accordance with a preferred embodiment of the present invention, the DNA transfer method is any suitable transfer method including DNA bombardment, electroporation, PEG-mediated DNA transfer and whiskers, among others.

30 In accordance with a preferred embodiment of the present invention, the expression construct comprises at least a nitrogen-inducible promoter and at least one cis- or trans-acting elements.

35 In accordance with a preferred embodiment of the present invention, the organism is a plant, a fungus, a bacteria, a yeast or an animal.

In accordance with a preferred embodiment of the present invention, the promoter or cis-acting sequence is isolated from the 5' upstream region of any gene involved in a nitrogen assimilatory pathway.

5 In accordance with a preferred embodiment of the present invention, the promoter or cis-acting sequence is isolated from the 5'upstream region of any gene for which transcription is modulated by availability of a given nitrogen source.

10 In accordance with a preferred embodiment of the present invention, the promoting or cis-acting sequence is any sequence for which transcriptional activity is regulated by addition or removal of any nitrogen source in or from any living organism's environment.

In accordance with a preferred embodiment of the present invention, the organism from which the promoter or cis-acting sequence is isolated from is any plant, fungus, yeast, bacteria or animal.

15 In accordance with a preferred embodiment of the present invention, there is provided a promoter for promoting transcription of a foreign gene in transgenic organisms, which comprises a nitrogen-inducible promoter with or without cis-acting sequence for expression of said gene and adapted to be modulated for transcriptional expression of
20 said gene by addition or removal of a nitrogen inducer.

Preferably, the promoter has a sequence selected from the group consisting of SEQ ID NOS: 1 to 13 and functional fragments and derivatives thereof.

25 In accordance with a preferred embodiment of the present invention, there is provided a terminator allowing expression of a foreign gene in transgenic organisms being used in combination with a promoter, which comprises a polyadenylation signal end site for insertion at the 3'end of said gene, wherein said terminator is operationally located with respect to said gene and said promoter and thereby allows expression of
30 said gene.

Preferably, the terminator has a sequence selected from the group consisting of SEQ ID NOS: 14 to 16 and functional fragments and derivatives thereof.

35 For the purpose of the present invention the following terms are defined below.

The expression "functional fragments or derivatives thereof" is intended to mean any derivative or fragment of sequences SEQ. ID. NOS:1-16 which allow for an equivalent level of expression of a foreign gene as the promoter of the present invention set forth in SEQ. ID. NOS:1-13 or as the terminator of the present invention set forth in SEQ. ID. NOS:14-16.

BRIEF DESCRIPTION OF THE DRAWINGS

Fig. 1 illustrates GUS expression level using promoter Nir (SEQ ID NOS:2-13) and terminator NOS in the leaves of transgenic tobacco plants before (empty columns) and after (filled columns) nitrate fertilization of the plants. Tobacco plants were transformed with full-length and deletions of the alfalfa NiR promoter and NOS terminator functionally positioned to control transcription and termination of the GUS reporter gene, as described in Methods. GUS activity was measured according to Jefferson et al. (1987, *EMBO J.* 13:3901-3907) prior and after induction by nitrate.

Fig. 2 illustrates GUS expression level using promoter Nir (SEQ ID NOS:2, 3, 5-6) and terminator Nir (SEQ ID NOS:15-16) in the leaves of transgenic tobacco plants before (empty columns) and after (filled columns) nitrate fertilization of the plants. Tobacco plants were transformed with full-length and deletions of the alfalfa NiR promoter and NiR terminator functionally positioned to control transcription and termination of the GUS reporter gene, as described in Materials and Methods below. GUS activity was measured according to Jefferson et al (1987, *EMBO J.* 13:3901-3907) prior and after induction by nitrate.

Fig. 3 illustrates GUS expression level using promoter NiR and terminator NOS in the leaves of nodulated transgenic alfalfa plants before (empty columns) and after (filled columns) nitrate fertilization. Alfalfa genotype 11.9 was transformed with constructs GC2-E, GC2-B, DC1-D and 35S functionally positioned to drive expression of reporter gene GUS, as described in Khoudi et al (1997, *Gene* 197:343-351). Following regeneration, transgenic plants were transferred to sterile vermiculite and inoculated with *Rhizobium* strain Balzac (Nitragin). Plants were allowed to grow for 3 weeks with repeated additions of nitrate-free Hoagland's

solution; GUS activity was then measured in first fully expanded leaves as described by Jefferson et al. (1987, *EMBO J.* 13:3901-3907). Plants were then fertilized with 40 mM nitrate for two days. GUS activity was again measured in first fully-expanded leaves. Data shown herein are ratios
5 between post- and pre-nitrate induction GUS specific activities.

DETAILED DESCRIPTION OF THE INVENTION

Following is a detailed description of the method used to generate transgenic tobacco and alfalfa lines that can be modulated in
10 their expression of a reporter gene. It should be remembered that variations could be brought to the method by which nitrogen-inducible promoters could be isolated, by which they could be linked to ORFs in the construct used for expression in plants, by which different cis- and trans-acting elements of the constructs are used and spatially arranged, by
15 which the inducibility by nitrogen is demonstrated and used, while remaining within the scope of this invention.

In this embodiment, a NiR cDNA strand was first isolated from alfalfa using RT-PCR with primers deduced from a consensus plant NiR sequence. This cDNA stretch was then used either to perform
20 upstream/downstream genome walking. The NiR promoter region and deletions, the 5'UTR and the NiR terminator were then positioned functionally to control transcription and terminaison of reporter gene GUS. These constructs were inserted into suitable expression vectors for DNA bombardment onto tobacco and alfalfa leaves and for *Agrobacterium*
25 mediated DNA transfer as described by Desgagnés et al. (1995, *Plant Cell Tissue Organ Cult.* 42:129-140). These two DNA transfer methods were used to demonstrate that expression of the reporter gene can be modulated by addition or removal of nitrate in the growing medium.

Materials and Methods

30

Biological material

E. coli strain DH5- α was used to perform all cloning steps. Cold resistant alfalfa genotype 11.9 was used for all experiments including

stable transformation using *A. tumefaciens* infection (Desgagnés et al. (1995, *Plant Cell Tissue Organ Cult.* 42:129-140).

Isolation of total RNA

5 Total RNA was extracted using a hot phenol method essentially as described by de Vries et al. (1988, B6 page 1, In: Gevin SB and Shilperoot RA editors, *Plant Molecular Biology Manual*, Dordrecht: Kluwer Academic Publisher).

10 RT-PCR

 RT-PCR was used to produce a DNA fragment corresponding to one abundant NiR mRNA molecular species from leaf total mRNA. A conserved region was first identified from 5 public plant NiR ORFs, namely Genbank sequences #AB006032 (*Arabidopsis* Nir mRNA), #
15 X66145 (*Tobacco* partial Nir mRNA), #U10419 (*Bean* complete Nir cds), #X07568 (*Spinach* Nir mRNA), and #U90429 (*Glycine* max Nir complete cds). Degenerated oligonucleotides were deduced from two conserved regions, namely Nir5 - 5' GATATTGATGTTAGACTCAAGTGGC 3' (SEQ ID NO:17), at the 5' end and Nir3 - 5' CACYSATTCCACTTCCTWGGC 3' (SEQ ID NO:18), at the 3' end of the coding strand. A reverse
20 transcription reaction was first performed with 200 units of M-MLV reverse transcriptase (RT) for 1 hour at 37°C using 1 µg of total leaf RNA, 4 mM dNTP (1 mM each), 5 µM random hexamer primers in a 1X M-MLV-RT buffer (50 mM Tris-HCl pH 8.3, 75 mM KCl, 3mM MgCl₂). The PCR
25 reaction was performed in a Perkin Elmer Cetus GenAmp PCR system 9600 (EG&G, Wellesley, MA), using 2.5 units of Taq DNA polymerase, 2 µM Nir5 primer, 2 µM Nir3 primer, 800 µM dNTPs (200 µM each) in a 1X PCR buffer (20 mM Tris-HCl pH 8.4, 50 mM KCl, 2 mM MgCl₂). The cycling program used was: an initial 4 min at 94°C, 30 cycles of 1 min at
30 94°C, 30 sec at 55°C, and 3 min at 72°C. An extension period of 7 min at 72°C was included in the program.

DNA sequencing

DNA sequencing was performed as described by Sanger et al. (1977, *P.N.A.S. USA*, 74:5643-5647).

5

Genome walking

Walking upstream of the Alfalfa Nir cDNA fragment cloned from the RT-PCR reaction was performed using the Universal Genome Walker Kit from Clontech Laboratories (Palo Alto, CA) (Cat. #1807-1). The NiR-specific custom primers used to amplify sequences upstream of the coding sequence were:

10

Nir1106r - 5' TTGTCACATCAGCACATCCGTCTTTGC 3' (SEQ ID NO:19), and

15

Nir1061r - 5' TCGCCAAGTATCTTGTTTGAGCACTTG 3' (SEQ ID NO:20).

The amplified 3775 bp fragment was subcloned into pGEM-T Easy vector (Promega, Madison, WI) (Cat #A1360) for further analysis. The resulting plasmid was named pGNir4c.

20

The downstream walking was performed as the upstream walking using the following NiR specific primers:

Nir1c - 5' ATGTCTTCCTTCTCAGTACGTTTCCTC 3' (SEQ ID NO:28), and

Nir138c - 5' CAAGTTGATGCATCAAGGTGGGAGCCTAGA 3' (SEQ ID NO:29).

25

The amplified 3508 bp fragment was subcloned into pGEM-T easy vector (Promega, Madison, WI) (Cat #A1360) for further analysis. The resulting plasmid was named pGN3'1.

Construction of expression cassettes and vectors

30

The cassettes for expression analysis using the GUS reporter gene were assembled as follows. A promoterless GUS cassette was digested from pBI101 with HindIII and EcoRI, and was inserted into the HindIII and EcoRI sites of the pUC19 polycloning site. The resulting plasmid was named pBI201 and was used for further constructs. The Nir upstream sequences were PCR amplified using the AP2 primer from the Universal Genome Walking Kit as upstream positioned primer, and either one of custom-designed downstream primers ending with a SmaI

35

restriction site. The 4 primers were positioned either in the 5' UTR region of the gene (Nir-23r-Sma-5'AGAGCCCCGGGAGAAGAGAGTGTGTTTG3' (SEQ ID NO:21)), at the end of the transit peptide coding sequence (Nir51r-Sma - 5' TTCTCCCGGGGGACGAGAGATGGATGGT 3' (SEQ ID NO:22)), 50 bp after the transit peptide coding sequence (Nir103r-Sma - 5' TTCTCCCGGGGTTGAA-ACAGGTGCAACTGA 3' (SEQ ID NO:23)), and 100 pb after the transit peptide coding sequence (Nir158r-Sma - 5' TTCTCCCGGGTAACCATCTTTTTCCTCA 3' (SEQ ID NO:24)) Amplification was performed under standard conditions with pGNir4c plasmid as template.

The amplified fragments were digested with specific restriction enzymes in order to produce 5' deletions of the Nir promoter. The pNir3k-23 was produced by digesting the fragment previously amplified by AP2 and Nir-23r-Sma primers with XmaI, and inserting the resulting fragment into pBI201 previously digested with XmaI. A similar strategy was used to produce the pNir3k51, pNir3k103, and pNir3k158 plasmids except that the downstream primers used were Nir51r-Sma, Nir103r-Sma, and Nir158r-Sma, respectively. The pNir2.2k-23 was produced from a SmaI-BglII digestion of the AP2 - Nir-23-Sma amplified fragment inserted into the pBI201 previously digested with SmaI and BamH1. The same strategy was used to produce the pNir2.2k51, pNir2.2k103, and pNir2.2k158 plasmids except that the downstream primers used were Nir51r-Sma, Nir103r-Sma, and Nir158r-Sma, respectively. Fidelity and orientation of the insertions were verified by digestion with restriction enzymes. These deletion fragments were ligated to the 5'terminus of the GUS reporter gene in pBI201, and used for transitory expression studies using DNA bombardment. Upon identification of the adequate deletion fragments, they were sub-cloned into a binary plant expression vector such as pBI101 (Clontech).

For the construction of the cassettes containing the NiR terminator downstream of the GUS gene in addition to the NiR promoter, the following NiR specific primers were used:
Nir2514c-Sac - 5' AGAAGAGCTCAGTATATAGGTATTTGGTGA 3'(SEQ ID NO:30)

Nir2728c-Sac - 5' AGAAGAGCTCTTGTACATTTGGATAAGTCA 3' (SEQ ID NO:31)

Nir3029r-Eco - 5' AGAAGAATTCGTTTTCCCGATACTTCAACT 3' (SEQ ID NO:32)

5 A 617 bp terminator fragment was PCR amplified using the primers Nir2514c-Sac and Nir3029r-Eco, and a 503 bp terminator fragment was PCR amplified using the primers Nir2728c-Sac and Nir3029r-Eco. The fragments obtained were digested with SacI and EcoRI and inserted into the plasmids containing the NiR-GUS constructs after
10 deletion of the NOS terminator between the SacI and EcoRI sites.

 These recombinant plasmids were used for stable integration through *A. tumefaciens* infection as described below.

15 ***Agrobacterium*-mediated DNA transfer and regeneration of transgenic alfalfa lines**

 The recombinant plasmids were introduced into *Agrobacterium tumefaciens* strain LBA4404 by electroporation as described in Khoudi et al (1997, *Gene* 197:343-351). Selected *Agrobacterium* strains were then
20 co-cultivated with leaf disks from genotype C5-1 for 4 days in the absence of selection pressure (kanamycin). Following this incubation period, leaf disks were washed and pampered, and then allowed to form calli onto medium B5H. Calli were then transferred for 21 days on SH medium for embryo induction and for 28 days on BOi2Y for embryo development. Torpedo-shaped embryos were removed from Boi2Y and placed on MS
25 medium for regeneration. Kanamycin was present in all cultivation medium except for co-cultivation and regeneration on MS. This method is described in length in Desgagnés et al (1995, *Plant Cell Tissue Organ Cult.* 42:129-140). Rooted plantlets were grown to maturity in the greenhouse. Integration of the transgene was verified by PCR
30 amplification of a NiR-GUS fragment from genomic DNA. The primers used were:

Nir-102c – 5' CACACTTCTTCACTCACCTCTCAA 3' (SEQ ID NO:25)

Nir-2016c – 5' ATCTAGGAGGGGCAGACATTG 3' (SEQ ID NO:26)

GUS228r – 5' TCGGTATAAAGACTTCGCGCTGAT 3' (SEQ ID NO:27)

35

***Agrobacterium*-mediated DNA transfer and regeneration of transgenic tobacco lines**

The recombinant plasmids were introduced into *Agrobacterium tumefaciens* strain LBA4404 by electroporation as described in Khoudi et al (1997, *Gene* 197:343-351). Selected strains were co-cultivated with leaf disks for 2 days on MS medium without kanamycin. After this period, the explants were transferred to the selection medium (MS with Kanamycin). The explants were kept on this medium for 3 weeks to allow the formation of calli and shoots from the transfected cells. The kanamycin resistant shoots were transferred into the rooting MS medium. Rooted plantlets were grown to maturity in the greenhouse. Integration of the transgene was verified by PCR amplification of a NiR-GUS fragment from genomic DNA. The primer used were:

- Nir-102c – 5' CACACTTCTTCACTCACCTCTCAA 3' (SEQ ID NO:25)
Nir-2016c – 5' ATCTAGGAGGGGCAGACATTG 3' (SEQ ID NO:26)
GUS228r – 5' TCGGTATAAAGACTTCGCGCTGAT 3' (SEQ ID NO:27)

Nitrate induction

Transgenic and non-transgenic tobacco and alfalfa plants were grown in vermiculite medium without nitrate. Mineral balance was kept by repeated additions of nitrate-free Hoagland's solutions (Hoagland and Varnon, 1950, Circular 347, *California Agr. Exp. Stat. Berkeley*). Nitrate induction was performed by watering the plants with 20-20-20 fertilizer at a concentration of 5gL⁻¹ or as an alternative with Hoagland's solution supplemented with 40 mM nitrate.

NiR promoter activity in tobacco leaves

The NiR derived promoters were placed upstream of the GUS reporter gene in transcriptional and translational fusions. The 5' deletions of the NiR promoter analyzed here consisted in (1) a putative full length promoter comprising 2813 bp upstream of the initial ATG of the coding region, (2) a 1905 bp version of the promoter, and (3) a shorter 1111 bp version of the promoter. The 3' end of the promoter was fused to the 5' end of the GUS coding region to form transcriptional and translational

fusions. Translational fusions analyzed allowed the production of β -glucuronidase containing (1) the NiR chloroplast transit peptide, (2) the NiR transit peptide with an additional 17 amino acids from NiR, and (3) the NiR transit peptide with an additional 36 amino acids from NiR. The twelve combinations of 5' and 3' deletions of the NiR promoter introduced into tobacco plants are presented in Fig. 1.

The gene constructs were transferred into tobacco plants using the *Agrobacterium*-mediated transfection method (Khouidi et al., 1997, *Gene* 197:343-351). Transgenic plants were transferred to growth chambers and analyzed for their leaf β -glucuronidase content before and after nitrate fertilization. Fig. 1 presents the median level of β -glucuronidase activity measured in the 1st expanded leaf of plantlets.

All the NiR derived promoters showed reactivity to nitrate induction. Between 5 and 10 fold increase of β -glucuronidase expression was generally observed, irrespectively of the promoter truncation, indicating that important nitrate responsive elements are contained within the first 1.1 kb upstream of the initial ATG. Both 5' and 3' deletions of the NiR promoter led to important modifications of β -glucuronidase activity. The highest level of GUS expression was obtained with the 2.8 kb promoter, indicating that the far upstream regions have a regulatory role for the level of NiR expression in the leaves.

The translational fusions of the promoter to the GUS coding region resulted in variable expression level depending on the extension of the 5' end of the promoter. However, the shortest fusion (containing the 17 a.a. NiR transit peptide fused to the amino-terminal end of the β -glucuronidase) constantly resulted in the highest level of activity for all three 5' end truncations. This short translational fusion, combined with the longest extension of upstream promoter regions gave rise to the strongest promoter (3kb+50). When induced, this specific construct resulted in more than 13-fold the level of GUS expression obtained with the 35S-GUS-NOS construct.

When considering the longest 5' extension of the promoter, the transcriptional fusion to the GUS gene (3kb-5) was ~7 times less effective than the short translational fusion (3kb+50). However, in it's induced state, the level of GUS expression in the plants harboring the 3kb-5 promoter deletion was more than 1,8-fold that observed in the 35S-GUS-NOS plants.

Taken together, the results presented here clearly indicate that the sequences upstream of the alfalfa NiR gene have the capacity to drive high and inducible expression of an exogenous gene in tobacco leaves.

10

Efficiency of NiR terminator

Tobacco plants were transformed with constructs consisting of promoter NiR and deletions (SEQ ID NOS: 2, 3, 5 and 6), and 35S, together with 3'UTR sequences and terminator (SEQ ID NOS: 15 and 16), functionally positioned to drive transcription and termination of reporter gene GUS. Growth, nitrate induction and GUS activity measurements were performed as per experiment illustrated in Fig 1. Results shown in Fig 2 demonstrate that the terminator sequence of Nir allows termination of transcription into a translatable messenger RNA.

20

NiR promoter activity in alfalfa leaves

Transgenic alfalfa plant containing the gene constructs presented in Fig. 3 were obtained using the *Agrobacterium*-mediated transfection method of Desgagnés et al. (1995, *Plant Cell Tissue Organ Cult.* 42:129-140). The *in vitro* plants were transferred into growth chamber to allow a normal vegetative growth. Cuttings from each transgenic line were grown in vermiculite and fertilized with nitrate-free Hoagland medium. After two weeks, the roots were inoculated with Nitragin (LiphaTech inc., Milwaukee, WI). Two weeks after inoculation, nodules had developed on the roots. Nodulated plants were allowed to continue their vegetative growth for at least another week before the fluorometric measurement of β -glucuronidase activity (before induction)

30

was performed. After the measurement, the plants were watered with Hoagland medium containing 40 mmol KNO₃. Two days after induction, leaf β -glucuronidase activity was measured to evaluate the nitrate inducibility of the NiR promoter in alfalfa leaves. Results are presented in

5 Fig. 3. Results show that promoter NiR induces expression of GUS reporter gene upon addition of nitrate in nodulated alfalfa plants. Taken together, this last series of result demonstrate that NiR promoter inducibility can be used to positively regulate expression of a foreign gene in alfalfa plants when fixation of atmospheric nitrogen is replaced by

10 nitrate assimilation.

While the invention has been described in connection with specific embodiments thereof, it will be understood that it is capable of further modifications and this application is intended to cover any variations, uses, or adaptations of the invention following, in general, the

15 principles of the invention and including such departures from the present disclosure as come within known or customary practice within the art to which the invention pertains and as may be applied to the essential features hereinbefore set forth, and as follows in the scope of the appended claims.

20

WHAT IS CLAIMED IS:


1. A method for regulating transcription of a foreign gene in transgenic organisms, comprising the steps of:
 - a) preparing a transgenic organism using an expression construct consisting of at least a nitrogen-inducible promoter having a sequence selected from the group consisting of SEQ ID NOS: 1 to 13 and functional fragments and derivatives thereof, and an ORF of a gene, wherein said promoter is operationally located with respect to said gene for expression of said gene.
2. The method of claim 1, which further comprises the step of regulating transcriptional expression of said gene by addition or removal of a nitrogen inducer.
3. A method for regulating transcription of a foreign gene in transgenic organisms comprising:
 - a) preparing an expression construct consisting of at least a nitrogen-inducible promoter with or without cis-acting sequence, an ORF of a gene, and a polyadenylation signal end site at the 3'end of said construct, wherein said promoter is operationally located with respect to said gene for expression of said gene and modulated for transcriptional expression of said gene by addition or removal of a nitrogen inducer;
 - b) sub-cloning the construct of step a) into a suitable transfection vector for said organism;
 - c) transferring said vector into DNA of said organism or a cell thereof; and
 - d) selecting for transgenicity on a suitable medium.
4. The method of claim 3, wherein said organism is a plant.
5. The method of claim 4, which further comprises the steps of:
 - e) introducing the vector into a suitable *Agrobacterium tumefaciens* strain;

- f) using the *Agrobacterium* strain of step a) to transfer T-DNA into a plant cell;
 - g) selecting for transgenicity of said plant cell on a suitable medium;
 - h) regenerating embryos or plantlets from said transgenic cells; and
 - i) growing mature plants from said regenerated embryos.
6. The method of claim 3, wherein the cis-acting sequence is isolated from 5' upstream region of an expressed Nir gene in alfalfa.
7. The method of claim 3, wherein the promoter has the sequence set forth in SEQ ID NOS:1 to 13 and functional fragments and derivatives thereof.
8. The method of claim 1, wherein the organism is selected from the group consisting of a plant, a fungus, a bacteria, a yeast and an animal.
9. The method of claim 8, wherein the plant is a dicot.
10. The method of claim 9, wherein the dicot is alfalfa or tobacco.
11. The method of claim 2, wherein the nitrogen inducer is nitrate.
12. The method of claim 3, wherein the DNA transfer method is any suitable transfer method including DNA bombardment, electroporation, PEG-mediated DNA transfer and whiskers.
13. The method of claim 3, wherein the expression construct comprises at least a nitrogen-inducible promoter and at least one cis- or trans-acting elements.


14. The method of claim 3, wherein the promoter or cis-acting sequence is isolated from the 5' upstream region of any gene involved in a nitrogen assimilatory pathway.
15. The method of claim 3, wherein the promoter or cis-acting sequence is isolated from the 5'upstream region of any gene for which transcription is modulated by availability of a nitrogen source.
16. The method of claim 3, wherein the promoting or cis-acting sequence is any sequence for which transcriptional activity is modulated by addition or removal of any nitrogen source in or from any living organism's environment.
17. The method of claim 3, wherein the organism from which the promoter or cis-acting sequence is isolated from is any plant, fungus, yeast, bacteria or animal.
18. A promoter for promoting transcription of a foreign gene in transgenic organisms, which comprises a nitrogen-inducible promoter with or without cis-acting sequence for expression of said gene and adapted to be modulated for transcriptional expression of said gene by addition or removal of a nitrogen inducer.
19. The promoter of claim 18, wherein said promoter has a sequence selected from the group consisting of SEQ ID NOS: 1 to 13 and functional fragments and derivatives thereof.
20. A terminator allowing expression of a foreign gene in transgenic organisms being used in combination with a promoter, which comprises a polyadenylation signal end site for insertion at the 3'end of said gene, wherein said terminator is operationally located with respect to said gene and said promoter and thereby allows expression of said gene.

21. The terminator of claim 19, wherein said terminator has a sequence selected from the group consisting of SEQ ID NOS: 14 to 16 and functional fragments and derivatives thereof.


1 / 3

*Fig. 1*

2 / 3

*Fig. 2*

3 / 3

*Fig. 3*

SEQUENCE LISTING

<110> VÉZINA, Louis-Philippe
D'AOUST, Marc-André
MEDICAGO INC.

<120> METHOD FOR REGULATING TRANSCRIPTION OF
FOREIGN GENES

<130> 14149-3 "PCT"

<150> US 60/157,133

<151> 1999-10-04

<160> 32

<170> FastSEQ for Windows Version 3.0

<210> 1

<211> 3714

<212> DNA

<213> Artificial Sequence

<220>

<223> Sequences to be used as promoter for regulating
expression of foreign genes

<400> 1

ctgtacattc	atcttgccgc	ctttgcattc	acttggccac	aaagagtaga	gagaaggaag	60
agaagagccc	agacttcaag	aagcgacctt	gcaagtgcac	tcgaggggtca	gaaactgtat	120
atcatatcta	tgtgagagaa	aggggaacat	ttgagatgga	gtccattttac	ttgaggtata	180
cttattatth	tgatcaataa	atttgtatac	ttcttattta	gatcaataaaa	tttgtcatta	240
agctataatc	caaaataaat	tacgatcaaa	tatgcaaagt	ttagccagta	cttgtgttaa	300
acttgatggc	atctcttggg	ttctttggca	atcacatgcc	taagaaataa	atagtatcat	360
atgatttgtg	ttggtcagac	ttcagagtca	gatgactctg	tttgataaaa	cagcttaatt	420
aagcgcttat	agaatatcat	atgatttgtg	ttggtcagac	ttcagagcat	ctcttgggtt	480
ctctggcaat	catatgccta	agaaataaat	agtatcatat	gatttgtgtt	ggtcagactt	540
cagagtcaga	tgacctgtt	tgggtaaaca	gcttaattaa	gtgcttatag	aataagcgct	600
tatcatataa	gtgcttttgt	acagttatth	ctatgaaagt	agaagaaata	gtcatattgt	660
tttaataata	gctatcctgg	agagcttgtg	gaaataacca	gaaaagaact	tatggacacg	720
tcatgagctg	tttacataag	atctccctaa	cagtctcaaa	agtgtttatg	ccagtagata	780
aattcaataa	agtcaatcta	aacagaccct	aaatccatta	tgggtacctat	catttttagct	840
tattccatct	ttattaagaa	tgtcatgaga	taacataatg	ataacacatt	atthttgacac	900
aaatgggcag	atctagcaat	ttaactctgg	agtccttcaa	gactgctgtt	cttacgaagt	960
tcacgtccct	gaatcatgtt	cctgtatgga	agcctgaaag	acctcaaatt	ctaaaagggtg	1020
gcgataaatt	gaaggttttac	aaaatatacc	ctgcgggctt	gacacagagg	caagctcttt	1080
ataccttcca	gttcaacggg	gatgttgatt	tcagaagtca	cttgagagac	aatccttgtg	1140
ccaagtttga	agtaatthtt	gtgtagcata	tgttgagcta	cctacaatth	acatgatcac	1200
ctagcattag	ctctttcact	taactgagag	aatgaagttt	taggaatgag	tatgaccatg	1260
gagtcggcat	ggctttgtaa	tgcctaccct	actttggcca	actcatcggg	gattttacatt	1320
cagaaaatat	acatgacttc	aaccataact	aaaccctttt	ttgtaagata	actgaatgtt	1380
catatthta	gttgggttgt	agtgtthttt	cttgattata	tccagacagt	tacaagttgg	1440
acaacaagat	tgtgggtctg	tactgttatt	tattttatth	tttttttagca	gaaacacctt	1500
atctthttgt	tcgtttgaat	gtagaatgaa	aataaaagaa	agaaaatata	acatcatcgg	1560
ccgcgcttgt	ctaatttcgg	gcagtttagga	tcctctccgg	tcaccggaaa	gtttcagtag	1620
aagaaacaaa	acaccgtgac	taaaatgata	ctattattht	atthattgtg	ttttctcttt	1680
ttctaccgga	actthtttaga	acggatccca	actcgttccg	gggccgctac	aactgaaaca	1740
aaagaagata	ttttctctct	cttcagaaat	gtaagthttt	ctttacagat	acctttcac	1800
catttgattc	agatgtgggtg	actagagata	aagcatacta	atthgactct	tggaaaccca	1860
taaagthtat	gttatccgtg	ttctggacca	atccacttgg	gggcataacc	tgtgtctatg	1920

tgtggtttgg	tttccattct	gatttatgcg	gcgacttgta	attttaaatac	taggagggggc	1980
agacattgaa	caatcccaat	attttaataa	cttatgcaag	atttttttta	ttaatgagat	2040
gatgtgtttg	tgactgagat	tgagtcatac	atttcactaa	gaaatgggtc	caagtaccaa	2100
actatcatga	cccagttgca	aacatgacgt	tcgggagtg	tcactttgat	agttcaattt	2160
catcttggct	tcttattcct	tttataattc	taattcttct	tgtgtaaact	atttcattga	2220
ttatttttct	ttaaaattta	catgtcattt	attttgcctc	actaactcaa	ttttgcatat	2280
aacaatgata	agtgatattt	tgactcacia	aatttacatc	aaatttcgac	atcgtttatt	2340
atgttcattg	gatgattaac	aaatataaca	aactttgcaa	ctaattaacc	accaactgaa	2400
tataattaac	tataactgtg	aaagtagtta	accatatttt	ttagatgtat	atatcatccg	2460
ttgaatgtaa	ttattcatat	atttgaacta	agttacccta	caacttaaa	aacttaaaaga	2520
actcggtttg	agacctgggg	acgaaaatgt	aatgagactt	taatgttgac	tttgacaccg	2580
caccacatgt	gccttttaca	tatagtttat	atgacaagta	atgacaatcc	ttgctctatt	2640
ataaggcgac	ccttagctcc	aaccaaagga	cgatggagtt	aagaaagaaa	ctcttgctta	2700
cttgtaagg	ccacacttct	tcactcacct	ctcaatttca	tcctacaaaa	atgtccaaac	2760
ttctctttct	cacaatcaca	aactcattcc	aaacacactc	tcttctccaa	aaatgtcttc	2820
cttctcagta	cgtttcctca	ccccaccatc	catctctcgt	ccaacaaaa	catggctact	2880
atctgctgca	actccatcag	ttgcacctgt	ttcaacacca	caagttgatg	catcaagggt	2940
ggagcctaga	gttgaggaaa	aagatgggta	ctgggttttg	aaggaagagt	atagaggggg	3000
tattaatcct	caggagaaa	ttaagattca	gaaagaacct	atgaagcttt	ttatggaagg	3060
tgggattaat	gatttgggta	atatgtctct	tgaagagatt	gaaagctcta	agcttactaa	3120
agatgatatt	gatgttagac	ttaaatgggt	tgggtctttt	catagaagga	aacatcattg	3180
taagtttttt	taccttcttt	ttatacctca	aagttctctc	atactctgta	tttgttttatt	3240
agtttttgta	gacttaaata	ttctctttga	tttacatagt	gaaactccat	tttgttttcc	3300
gaaattgtag	tgtgtatagt	ctagaaaatt	aagaagtaga	caaaatgatt	tatgagattg	3360
taaattgtag	gctttttatc	aattttattaa	tttttagagac	caaaatttgc	ctatcttatt	3420
tggaccaata	ttgtatgtca	ggatcgacat	gagtttagta	aaatcatgac	ggcaccatga	3480
ctgtgttgaa	gcttctttgt	gtaactttta	ccaaaattat	atggcacacc	ataattatgc	3540
aaactcacgg	tcgatccaaa	catagaaatt	cgggtgtaat	ctttgtgaga	ataaaaagct	3600
atgagttatg	ttgtactaat	ttatttccat	tgtgaaaatc	agatggtaga	tttatgatga	3660
gactgaaact	tccaaatggg	gtaacaacaa	gtgctcaaac	aagatacttg	gcga	3714

<210> 2

<211> 2808

<212> DNA

<213> Artificial Sequence

<220>

<223> Sequences to be used as promoter for regulating expression of foreign genes

<400> 2

ctgtacattc	atcttgccgc	ctttgcattc	acttggccac	aaagagtaga	gagaaggaag	60
agaagagccc	agacttcaag	aagcgacctt	gcaagtgcac	tcgaggggtca	gaaactgtat	120
atcatatcta	tgtgagagaa	aggggaacat	ttgagatgga	gtccattttac	ttgaggtata	180
cttattattt	tgatcaataa	atttgtatac	ttcttattta	gatcaataaa	tttgtcatta	240
agctataatc	caaaataaat	tacgatcaaa	tatgcaaatg	ttagccagta	cttgtgttaa	300
acttgatggc	atctcttggg	ttctttggca	atcacatgcc	taagaaataa	atagtatcat	360
atgatttgtg	ttgggtcagac	ttcagagtca	gatgactctg	tttggataaa	cagcttaatt	420
aagcgcttat	agaatatcat	atgatttgtg	ttgggtcagac	ttcagagcat	ctcttgggtt	480
ctctggcaat	catatgccta	agaaataaat	agtatcatat	gatttgtgtt	ggtcagactt	540
cagagtcaga	tgaccctgtt	tgggtaaaca	gcttaattaa	gtgcttatag	aataagcgct	600
tatcatataa	gtgcttttgt	acagttattt	ctatgaaagt	agaagaaata	gtcatattgt	660
tttaatatata	gctatcctgg	agagcttgtg	gaaataacca	gaaaagaact	tatggacacg	720
tcattgagctg	tttacataag	atctccctaa	cagctcctaa	agtgtttatg	ccagtagata	780
aattcaataa	agtcaatcta	aacagaccct	aatccatta	tggtacctat	catttttagct	840
tattccatct	ttattaagaa	tgtcatgaga	taacataatg	ataacacatt	attttgacac	900
aatggggcag	atctagcaat	ttaactctgg	agtccttcaa	gactgctgtt	cttacgaagt	960
tcacgtccct	gaatcatgtt	cctgtatgga	agcctgaaag	acctcaaatt	ctaaaagggtg	1020
gcgataaatt	gaagggtttac	aaaatatacc	ctgcgggctt	gacacagagg	caagctcttt	1080
ataccttcca	gttcaacggg	gatgttgatt	tcagaagtca	cttggagagc	aatccttgtg	1140

ccaagtttga	agtaattttt	gtgtagcata	tgttgagcta	cctacaattt	acatgatcac	1200
ctagcattag	ctctttcact	taactgagag	aatgaagttt	taggaatgag	tatgaccatg	1260
gagtcggcat	ggctttgtaa	tgcctaccct	actttggcca	actcatcggg	gatttacatt	1320
cagaaaatat	acatgacttc	aaccatactt	aaaccctttt	ttgtaagata	actgaatggt	1380
catatttaat	gttgggttgt	agtgttttta	cttgattata	tccagacagt	tacaagttgg	1440
acaacaagat	tgtgggtctg	tactgttatt	tatttttttt	tttttttagca	gaaacacctt	1500
atctttttgt	tcgtttgaat	gtagaatgaa	aataaaaagaa	agaaaatata	acatcatcgg	1560
ccgcgcctgt	ctaatttcgg	gcagtttaga	tcctctccgg	tcaccggaaa	gtttcagtag	1620
aagaaaacaaa	acaccgtgac	taaaatgata	ctattatttt	atttattgtg	tttttctttt	1680
ttctaccgga	acttttttga	acggatccca	actcgtttccg	gggcccgtac	aactgaaaca	1740
aaagaagata	ttttctctct	cttcagaaat	gtaagttttc	ctttacagat	accttaccac	1800
catttgattc	agatgtggtg	actagagata	aagcactacta	atttgactct	tggaaaccca	1860
taaagtttat	gttatccgtg	ttctggacca	atccacttgg	gggcataacc	tgtgtctatg	1920
tgtggttttg	tttccattct	gatttatgcg	gcgacttgta	atttaaaatc	taggaggggc	1980
agacattgaa	caatcccaat	attttaataa	cttatgcaag	atttttttta	ttaatgagat	2040
gatgtgtttg	tgactgagat	tgagtcatac	atttcactaa	gaaatgggtc	caagtaccaa	2100
actatcatga	cccagttgca	aacatgacgt	tcgggagtg	tcactttgat	agttcaattt	2160
catcttggtc	tcttattcct	tttataattc	taattcttct	tgtgtaaact	atttcatgta	2220
ttatttttct	ttaaaattta	catgtcattt	attttgcctc	actaactcaa	ttttgcatat	2280
aacaatgata	agtgatattt	tgactcacia	aatttacatc	aaatttcgac	atcgtttatt	2340
atgttcattg	gatgattaac	aaatataaca	aactttgcaa	ctaattaacc	accaactgaa	2400
tataattaac	tataactgtg	aaagtagtta	accatatttt	ttagatgtat	atatcatccg	2460
ttgaatgtaa	ttattcatat	atttgaacta	agttacccta	caacttaag	aacttaagaa	2520
actcggtttg	agacctgggg	acgaaaatgt	aatgagactt	taatgttgac	tttgacaccg	2580
caccacatgt	gcctttttaca	tatagtttat	atgacaagta	atgacaatcc	ttgctctatt	2640
ataaggcgac	ccttagctcc	aaccaaagga	cgatggagtt	aagaaagaaa	ctcttgctta	2700
cttgtgaagg	ccacacttct	tcactcacct	ctcaatttca	tcctacaaaa	atgtccaaac	2760
ttctctttct	cacaatcaca	aactcattcc	aaacacactc	tcttctcc		2808

<210> 3

<211> 2069

<212> DNA

<213> Artificial Sequence

<220>

<223> Sequences to be used as promoter for regulating expression of foreign genes

<400> 3

gatctcccta	acagtctcaa	aagtgtttat	gccagtagat	aaattcaa	aagtcaatct	60
aaacagaccc	taaatccatt	atggtacctt	tcatttttagc	ttattccatc	tttattaaga	120
atgtcatgag	ataacataat	gataacacat	tatttttgaca	caaatgggca	gatctagcaa	180
tttaactctg	gagtccttca	agactgctgt	tcttacgaag	ttcacgtccc	tgaatcatgt	240
tctgtatg	aagcctgaaa	gacctcaaat	tctaaaagg	ggcgataaat	tgaaggttta	300
caaaatatac	cctgcgggct	tgacacagag	gcaagctctt	tataccttcc	agttcaacgg	360
ggatgttgat	ttcagaagtc	acttgagag	caatccttgt	gccaagtttg	aagtaatttt	420
tgtgtagcat	atgttgagct	acctacaatt	tacatgatca	cctagcatta	gctctttcac	480
ttactgaga	gaatgaagtt	ttaggaatga	gtatgaccat	ggagtcggca	tggctttgta	540
atgcctaccc	tactttggcc	aactcatcgg	ggattttacat	tcagaaaata	tacatgactt	600
caaccatact	taaacccttt	tttgtaagat	aactgaatgt	tcatatttaa	tgttgggttg	660
tagtggtttt	acttgattat	atccagacag	ttacaagttg	gacaacaaga	ttgtgggtct	720
gtactgttat	ttattttatt	tttttttagc	agaaacacct	tatcttttgt	ttcgtttgaa	780
tgtagaatga	aaataaaaaga	aagaaaatat	aacatcatcg	gccgcgcttg	tctaatttcg	840
ggcagttagg	atcctctccg	gtcaccggaa	agtttcagta	gaagaaacaa	aacaccgtga	900
ctaaaatgat	actattattt	tattttattgt	gtttttcttt	tttctaccgg	aacttttttag	960
aacggatccc	aactcgttcc	ggggccgcta	caactgaaac	aaaagaagat	attttctctc	1020
tcttcagaaa	tgtaagtttt	cctttacaga	taccattcca	ccatttgatt	cagatgtggt	1080
gactagagat	aaagcatact	aatttgactc	ttggaaaacc	ataaagttta	tgttatccgt	1140
gttctggacc	aatccacttg	ggggcataac	ctgtgtctat	gtgtggtttg	gtttccattc	1200
tgatttatgc	ggcgacttgc	aattttaa	ctaggagggg	cagacattga	acaatcccaa	1260

tattttaata	acttatgcaa	gatttttttt	attaatgaga	tgatgtgttt	gtgactgaga	1320
ttgagtcata	catttcacta	agaaatggtt	ccaagtacca	aactatcatg	acccagttgc	1380
aaacatgacg	ttcgggagtg	gtcactttga	tagttcaatt	tcattcttggc	ttctttattcc	1440
ttttataatt	ctaattcttc	ttgtgtaaac	tattttcatgt	attatttttc	tttaaaattt	1500
acatgtcatt	tattttgcct	cactaactca	atttttgcata	taacaatgat	aagtgatatt	1560
ttgactcaca	aaatttacat	caaatttcga	catcgtttat	tatgttcatt	ggatgattaa	1620
caaataatac	aaactttgca	actaattaac	caccaactga	atataattaa	ctataactgt	1680
gaaagtagtt	aaccatattt	tttagatgta	tatatcatcc	gttgaatgta	attattcata	1740
tatttgaact	aagttaccct	acaacttaaa	gaacttaaa	aactcggttt	gagacctggg	1800
gacgaaaatg	taatgagact	ttaatgttga	ctttgacacc	gcaccacatg	tgcccttttac	1860
atatagttta	tatgacaagt	aatgacaatc	cttgctctat	tataaggcga	cccttagctc	1920
caaccaaagg	acgatggagt	taagaaagaa	actcttgctt	acttgtaagg	tccacacttc	1980
ttcactcacc	tctcaatttc	atcctacaaa	aatgtccaaa	cttctctttc	tcacaatcac	2040
aaactcattc	caaacacact	ctcttctcc				2069

<210> 4

<211> 1220

<212> DNA

<213> Artificial Sequence

<220>

<223> Sequences to be used as promoter for regulating expression of foreign genes

<400> 4

gacctctcc	ggtcaccgga	aagtttccagt	agaagaaaca	aaacaccgtg	actaaaatga	60
tactattatt	ttattttattg	tgtttttctt	ttttctaccg	gaacttttta	gaacggatcc	120
caactcgttc	cggggccgct	acaactgaaa	caaaagaaga	tattttctct	ctcttcagaa	180
atgtaagt	tcctttacag	atacccttc	accatttgat	tcagatgtgg	tgactagaga	240
taaagcatat	taatttgact	cttggaacc	cataaagttt	atgttatccg	tggtctggac	300
caatccactt	gggggcataa	cctgtgtcta	tgtgtgggtt	ggtttccatt	ctgatttatg	360
cggcgacttg	taatttataa	tctaggagg	gcagacattg	aacaatccca	atattttaat	420
aacttatgca	agattttttt	tattaatgag	atgatgtgtt	tgtgactgag	attgagtcac	480
acatttcact	aagaaatggt	tccaagtacc	aaactatcat	gacctcagttg	caaacatgac	540
gttcgggagt	ggtcactttg	atagttcaat	ttcatcttgg	cttcttattc	cttttataat	600
tctaattctt	cttgtgtaaa	ctatttcatg	tattattttt	ctttaaaatt	tacatgtcat	660
ttattttgcc	tcactaactc	aattttgcat	ataacaatga	taagtgatat	tttgactcac	720
aaaattttaca	tcaaatttcg	acatcgttta	ttatgttcat	tggatgatta	acaaatataa	780
caaactttgc	aactaattaa	ccaccaactg	aatataatta	actataactg	tgaaagtagt	840
taaccatatt	tttttagatgt	atatatcatc	cgttgaatgt	aattattcat	atatttgaac	900
taagttaccc	tacaacttaa	agaacttaaa	gaactcgggt	tgagacctgg	ggacgaaaat	960
gtaatgagac	tttaatgttg	actttgacac	cgcaccacat	gtgcctttta	catatagttt	1020
atatgacaag	taatgacaat	ccttgctcta	ttataaggcg	acccttagct	ccaaccaaag	1080
gacgatggag	ttaagaaaga	aactcttgct	tacttgtaag	gtccacactt	cttcactcac	1140
ctctcaattt	catcctacaa	aaatgtccaa	acttctcttt	ctcacaatca	caaactcatt	1200
ccaaacacac	tctcttctcc					1220

<210> 5

<211> 2863

<212> DNA

<213> Artificial Sequence

<220>

<223> Sequences to be used as promoter for regulating expression of foreign genes

<400> 5

ctgtacattc	atcttgccgc	ctttgcattc	acttggccac	aaagagtaga	gagaagggaag	60
agaagagccc	agacttcaag	aagcgacctt	gcaagtgcac	tgcaggggtca	gaaactgtat	120
atcatatcta	tgtgagagaa	aggggaacat	ttgagatgga	gtccatttac	ttgaggtata	180

cttattattt	tgatcaataa	atttgtatac	ttcttattta	gatcaataaa	tttgtcatta	240
agctataatc	caaaaataaat	tacgatcaaa	tatgcaaagt	ttagccagta	cttgtgttaa	300
acttgatggc	atctcttggg	ttctttggca	atcacatgcc	taagaaataa	atagtatcat	360
atgatttgtg	ttggtcagac	ttcagagtca	gatgactctg	tttggataaa	cagcttaatt	420
aagcgcttat	agaatatcat	atgatttgtg	ttggtcagac	ttcagagcat	ctcttgggtt	480
ctctggcaat	catatgccta	agaaataaat	agtatcatat	gatttgtgtt	ggtcagacct	540
cagagtcaga	tgaccctgtt	tgggtaaaca	gcttaattaa	gtgcttatag	aataagcgct	600
tatcatataa	gtgcttttgt	acagttattt	ctatgaaaag	agaagaaata	gtcatattgt	660
tttaataata	gctatcctgg	agagcttgtg	gaaataacca	gaaaagaact	tatggacacg	720
tcatgagctg	tttacataag	atctccctaa	cagtctcaaa	agtgtttatg	ccagtagata	780
aattcaaata	agtcaatcta	aacagaccct	aaatccatta	tggtagctat	catttttagct	840
tattccatct	ttattaagaa	tgtcatgaga	taacataatg	ataacacatt	attttgacac	900
aaatgggcag	atctagcaat	tttaactctg	agtccttcaa	gactgctgtt	cttacgaagt	960
tcacgtccct	gaatcatgtt	cctgtatgga	agcctgaaaag	acctcaaatt	ctaaaagggtg	1020
gcgataaatt	gaagggtttac	aaaatatacc	ctgcgggctt	gacacagagg	caagctcttt	1080
ataccttcca	gttcaacggg	gatgttgatt	tcagaagtca	cttggagagc	aatccttgtg	1140
ccaagtttga	agtaattttt	gtgtagcata	tgttgagcta	cctacaattt	acatgatcac	1200
ctagcattag	ctctttcact	taactgagag	aatgaagttt	taggaatgag	tatgaccatg	1260
gagtcggcat	ggctttgtaa	tgcctaccct	actttggcca	actcatcggg	gatttacatt	1320
cagaaaatat	acatgacttc	aaccatactt	aaaccccttt	ttgtaagata	actgaatgtt	1380
catattttaa	gttgggttgt	agtgttttta	cttgattata	tccagacagt	tacaagttgg	1440
acaacaagat	tgtgggtctg	tactgttatt	tatttatttt	tttttttagca	gaaacacctt	1500
atcttttgtt	tcgtttgaat	gtagaatgaa	aataaaaagaa	agaaaatata	acatcatcgg	1560
ccgcgcttgt	ctaatttcgg	gcagtttagga	tcctctccgg	tcaccggaaa	gtttcagtag	1620
aagaaacaaa	acaccgtgac	taaaatgata	ctattatttt	atttattgtg	ttttctcttt	1680
ttctaccgga	acttttttaga	acggatccca	actcgttccg	gggccgctac	aactgaaaca	1740
aaagaagata	ttttctctct	cttcagaaat	gtaagttttc	ctttacagat	accttccac	1800
catttgattc	agatgtgggtg	actagagata	aagcatacta	atttgactct	tggaaaccca	1860
taaagtttat	gttatccgtg	ttctggacca	atccacttgg	gggcataacc	tgtgtctatg	1920
tgtggtttgg	tttccattct	gattttatg	gcgacttgta	attttaaatac	taggaggggc	1980
agacattgaa	caatcccaat	attttaataa	cttatgcaag	atttttttta	ttaatgagat	2040
gatgtgtttg	tgactgagat	tgagtcatac	atttcactaa	gaaatgggtc	caagtaccaa	2100
actatcatga	cccagttgca	aacatgacgt	tcgggagtg	tcactttgat	agttcaattt	2160
catcttggct	tcttattcct	tttataattc	taattcttct	tgtgtaaaact	atttcatgta	2220
ttatttttct	ttaaaattta	catgtcattt	attttgctc	actaactcaa	ttttgcatat	2280
aacaatgata	agtgatattt	tgactcacia	aattttacatc	aaatttcgac	atcgtttatt	2340
atgttcattg	gatgattaac	aaatataaca	aactttgcaa	ctaattaacc	accaactgaa	2400
tataattaac	tataactgtg	aaagtagtta	accatatttt	ttagatgtat	atatcatccg	2460
ttgaatgtaa	ttattcatat	atttgaacta	agttacccta	caacttaaag	aacttaaaga	2520
actcggtttg	agacctgggg	acgaaaatgt	aatgagactt	taatgttgac	tttgacaccg	2580
caccacatgt	gcctttttaca	tatagtttat	atgacaagta	atgacaatcc	ttgctctatt	2640
ataaggcgac	ccttagctcc	aaccaaagga	cgatggagtt	aagaaagaaa	ctcttgctta	2700
cttgaagggt	ccacacttct	tcactcacct	ctcaatttca	tcctacaaaa	atgtccaaac	2760
ttctctttct	cacaatcaca	aactcattcc	aaacacactc	tcttctccaa	aaatgtcttc	2820
cttctcagta	cgtttctctca	cccaccatc	catctctcgt	ccc		2863

<210> 6

<211> 2124

<212> DNA

<213> Artificial Sequence

<220>

<223> Sequences to be used as promoter for regulating expression of foreign genes

<400> 6

gatctcccta	acagtctcaa	aagtgtttat	gccagtagat	aaattcaaat	aagtcaatct	60
aaacagaccc	taaatccatt	atggtagcta	tcattttagc	ttattccatc	tttattaaga	120
atgtcatgag	ataacataat	gataacacat	tattttgaca	caaatgggca	gatctagcaa	180
tttaactctg	gagtccttca	agactgctgt	tcttacgaag	ttcacgtccc	tgaatcatgt	240

tcctgtatgg	aagcctgaaa	gacctcaaat	tctaaaaggt	ggcgataaat	tgaaggttta	300
caaaatatac	cctgcgggct	tgacacagag	gcaagctctt	tataccttcc	agttcaacgg	360
ggatgttgat	ttcagaagtc	acttggagag	caatccttgt	gccagtttg	aagtaatttt	420
tgtgtagcat	atgttgagct	acctacaatt	tacatgatca	cctagcatta	gctctttcac	480
ttaactgaga	gaatgaagtt	ttaggaatga	gtatgaccat	ggagtcggca	tggctttgta	540
atgcctaccc	tactttggcc	aactcatcgg	ggattttacat	tcagaaaata	tacatgactt	600
caaccatact	taaacccctt	tttgtaagat	aactgaatgt	tcataattta	tgttggggtg	660
tagtggtttt	acttgattat	atccagacag	ttacaagttg	gacaacaaga	ttgtgggtct	720
gtactgttat	ttattttatt	tttttttagc	agaaacacct	tatcttttgt	ttcgtttgaa	780
tgtagaatga	aaataaaaaga	aagaaaatat	aacatcatcg	gccgcgcttg	tctaatttcg	840
ggcagttagg	atcctctccg	gtcaccggaa	agtttcagta	gaagaaacaa	aacaccgtga	900
ctaaaatgat	actatttatt	tattttattgt	gtttttcttt	tttctaccgg	aacttttttag	960
aacggatccc	aactcgttcc	ggggccgcta	caactgaaac	aaaagaagat	attttctctc	1020
tcttcagaaa	tgttaagttt	cctttacaga	taccatttca	ccatttgatt	cagatgtggg	1080
gactagagat	aaagcatact	aatttgactc	ttggaaaccc	ataaagttta	tgttatccgt	1140
gttctggacc	aatccacttg	ggggcataac	ctgtgtctat	gtgtgggttg	gtttccattc	1200
tgattttatgc	ggcgacttgt	aattttaa	ctaggagggg	cagacattga	acaatcccaa	1260
tatttttaata	acttatgcaa	gatttttttt	attaatgaga	tgatgtgttt	gtgactgaga	1320
ttgagtcata	catttcacta	agaaatgggt	ccaagtacca	aactatcatg	accagttgc	1380
aaacatgacg	ttcgggagtg	gtcactttga	tagttcaatt	tcactctggc	ttcttattcc	1440
ttttataatt	ctaattcttc	ttgtgtaa	tatttcatgt	attatttttc	tttaaaattt	1500
acatgtcatt	tattttgcct	cactaactca	attttgcata	taacaatgat	aagtgatatt	1560
ttgactcaca	aaattttacat	caaatttcga	catcgtttat	tatgttcatt	ggatgattaa	1620
caaatataac	aaactttgca	actaattaac	caccaactga	atataattaa	ctataactgt	1680
gaaagtagtt	aaccatattt	tttagatgta	tatatcatcc	gttgaatgta	attattcata	1740
tatttgaact	aagttaccct	acaacttaaa	gaacttaaa	aactcgggtt	gagacctggg	1800
gacgaaaatg	taatgagact	ttaatgttga	ctttgacacc	gcaccacatg	tgccctttac	1860
atatagttaa	tatgacaagt	aatgacaatc	cttgctctat	tataaggcga	cccttagctc	1920
caaccaaagg	acgatggagt	taagaaagaa	actcttgctt	acttgtaagg	tccacacttc	1980
ttcactcacc	tctcaatttc	atcctacaaa	aatgtccaaa	cttctctttc	tcacaatcac	2040
aaactcattc	caaacacact	ctcttctcca	aaaatgtctt	ccttctcagt	acgtttcctc	2100
acccaccat	ccatctctcg	tccc				2124

<210> 7

<211> 1160

<212> DNA

<213> Artificial Sequence

<220>

<223> Sequences to be used as promoter for regulating expression of foreign genes

<400> 7

gatcccaact	cgttccgggg	ccgtacaac	tgaacaaaa	gaagatattt	tctctctctt	60
cagaaatgta	agttttcctt	tacagatacc	cattcaccat	ttgattcaga	tgtgggtgact	120
agagataaag	catactaatt	tgactccttg	aaaccataa	agtttatgtt	atccgtgttc	180
tggaccaatc	cacttggggg	cataacctgt	gtctatgtgt	ggtttgggtt	ccattctgat	240
ttatgcggcg	acttgtaatt	taaaatctag	gaggggcaga	cattgaacaa	tcccaatatt	300
ttaataactt	atgcaagatt	ttttttatta	atgagatgat	gtgtttgtga	ctgagattga	360
gtcatacatt	tcactaagaa	atggttccaa	gtaccaaact	atcatgaccc	agttgcaaac	420
atgacgttcg	ggagtgggtca	ctttgatagt	tcaatttcat	cttggcttct	tattcctttt	480
ataattctaa	ttcttcttgt	gtaaactatt	tcattgtatta	tttttcttta	aaatttacat	540
gtcattttatt	ttgcctcact	aactcaattt	tgcatataac	aatgataagt	gatattttga	600
ctcacaaaat	ttacatcaaa	tttcgacatc	gtttattatg	ttcattggat	gattaacaaa	660
tataacaaac	tttgcaacta	attaaccacc	aactgaatat	aattaactat	aactgtgaaa	720
gtagttaacc	atatttttta	gatgtatata	tcattccgttg	aatgtaatta	ttcatatatt	780
tgaactaagt	taccctacaa	cttaagaac	ttaaagaact	cggtttgaga	cctggggacg	840
aaaatgtaat	gagactttta	tgttgacttt	gacaccgcac	cacatgtgcc	ttttacatat	900
agttttatatg	acaagtaatg	acaatccttg	ctctattata	aggcgaccct	tagctccaac	960
caaaggacga	tggagttaag	aaagaaactc	ttgcttactt	gtaagggtcca	cacttcttca	1020

ctcacctctc	aatttcatcc	tacaaaaatg	tccaaacttc	tctttctcac	aatcacaaac	1080
tcattccaaa	cacactctct	tctccaaaaa	tgtcttcctt	ctcagtagct	ttcttcaccc	1140
caccatccat	ctctcgtccc					1160

<210> 8

<211> 2904

<212> DNA

<213> Artificial Sequence

<220>

<223> Sequences to be used as promoter for regulating expression of foreign genes

<400> 8

ctgtacattc	atcttgccgc	ctttgcattc	acttggccac	aaagagtaga	gagaaggaag	60
agaagagccc	agacttcaag	aagcgacctt	gcaagtgcac	tcgagggtca	gaaactgtat	120
atcatatcta	tgtgagagaa	aggggaacat	ttgagatgga	gtccatttac	ttgagggtata	180
cttattatct	tgatcaataa	atctgtatac	ttcttattta	gatcaataaa	tttgtcatta	240
agctataatc	caaaataaat	tacgatcaaa	tatgcaaagt	ttagccagta	cttgtgttaa	300
acttgatggc	atctcttggg	ttctttggca	atcacatgcc	taagaaataa	atagtagcat	360
atgatttgtg	ttgggtcagac	ttcagagtca	gatgactctg	tttggataaa	cagcttaatt	420
aagcgcttat	agaatatcat	atgatttgtg	ttgggtcagac	ttcagagcat	ctcttgggtt	480
ctctggcaat	catatgccta	agaaataaat	agtatcatat	gatttgtgtt	gggtcagactt	540
cagagtcaga	tgacctgtgt	tgggtaaaca	gcttaattaa	gtgcttatag	aataagcgct	600
tatcatataa	gtgcttttgt	acagttatct	ctatgaaagt	agaagaaata	gtcatattgt	660
tttaataata	gctatcctgg	agagcttgtg	gaaataacca	gaaaagaact	tatggacacg	720
tcattgagctg	tttacataag	atctccctaa	cagtctcaaa	agtgtttatg	ccagtagata	780
aattcaataa	agtcaatcta	aacagaccct	aaatccatta	tggtacctat	catttttagct	840
tattccatct	ttattaagaa	tgtcatgaga	taacataatg	ataacacatt	attttgacac	900
aaatgggcag	atctagcaat	ttactctggg	agtccttcaa	gactgctgtt	cttacgaagt	960
tcacgtccct	gaatcatgtt	cctgtatgga	agcctgaaaag	acctcaaatt	ctaaaagggtg	1020
gcgataaatt	gaagggtttac	aaaatatacc	ctgcggggctt	gacacagagg	caagctcttt	1080
ataccttcca	gttcaacggg	gatgttgatt	tcagaagtca	cttggagagc	aatccttgtg	1140
ccaagtttga	agtaattttt	gtgtagcata	tggtgagcta	cctacaattt	acatgatcac	1200
ctagcattag	ctcttttact	taactgagag	aatgaagttt	taggaatgag	tatgaccatg	1260
gagtcggcat	ggctttgtta	tgctaccct	actttggcca	actcatcggt	gatttacatt	1320
cagaaaatat	acatgacttc	aaccatactt	aaaccccttt	ttgtaagata	actgaatgtt	1380
catattttaat	gttgggttgt	agtgttttta	cttgattata	tccagacagt	tacaagttgg	1440
acaacaagat	tgtgggtctg	tactgttatt	tattttattt	tttttttagca	gaaacacctt	1500
atcttttgtt	tcgtttgaat	gtagaatgaa	aataaaagaa	agaaaatata	acatcatcgg	1560
ccgcgcttgt	ctaatttcgg	gcagtttaga	tcctctccgg	tcaccggaaa	gtttcagtag	1620
aagaaacaaa	acaccgtgac	taaaatgata	ctattatttt	atttatttgt	tttttctttt	1680
ttctaccgga	actttttaga	acggatccca	actcgttccg	gggcccgtac	aactgaaaca	1740
aaagaagata	ttttctctct	cttcagaaat	gtaagttttc	ctttacagat	accttccac	1800
catttgattc	agatgtgggt	actagagata	aagcatacta	atttgactct	tggaaacca	1860
taaaagtttat	gttatccgtg	ttctggacca	atccacttgg	gggcataacc	tgtgtctatg	1920
tgtgggttgg	tttccattct	gatttatcgg	gcgacttgta	attttaaata	taggaggggc	1980
agacattgaa	caatcccaat	attttaataa	cttatgcaag	atttttttta	ttaatgagat	2040
gatgtgtttg	tgactgagat	tgagtcatac	atttcactaa	gaaatgggtc	caagtaccaa	2100
actatcatga	cccagttgca	aacatgacgt	tcgggagtggt	tcactttgat	agttcaattt	2160
catcttggct	tcttattcct	tttataattc	taattcttct	tgtgtaaact	atttcattga	2220
ttatttttct	ttaaaattta	catgtcattt	attttgcctc	actaactcaa	ttttgcatat	2280
aacaatgata	agtgatattt	tgactcacia	aattttacatc	aaatttcgac	atcgtttatt	2340
atgttcattg	gatgattaac	aaatataaca	aactttgcaa	ctaattaacc	accaactgaa	2400
tataattaac	tataactgtg	aaagtagtta	accatatttt	ttagatgtat	atatcatccg	2460
ttgaatgtaa	ttattcatat	atttgaacta	agttacccta	caacttaaag	aacttaaaga	2520
actcggtttg	agacctgggg	acgaaaatgt	aatgagactt	caatgttgac	tttgacaccg	2580
caccacatgt	gccttttaca	tatagtttat	atgacaagta	atgacaatcc	ttgtcttatt	2640
ataaggcgac	ccttagctcc	aaccaaagga	cgatggaggt	aagaaagaaa	ctcttgctta	2700
cttgtaagggt	ccacacttct	tcactcacct	ctcaatttca	tcctacaaaa	atgtccaaac	2760

ttctctttct	cacaatcaca	aactcattcc	aaacacactc	tcttctccaa	aaatgtcttc	2820
cttctcagta	cgtttctctca	ccccaccatc	catctctcgt	cccaacaaaa	catggctact	2880
atctgctgca	actccatcag	ttgc				2904

<210> 9

<211> 2165

<212> DNA

<213> Artificial Sequence

<220>

<223> Sequences to be used as promoter for regulating
expression of foreign genes

<400> 9

gatctcccta	acagtctcaa	aagtgtttat	gccagtagat	aaattcaaat	aagtcaatct	60
aaacagaccc	taaatccatt	atggtaccta	tcatttttagc	ttattccatc	tttattaaga	120
atgtcatgag	ataacataat	gataacacat	tatttttgaca	caaatgggca	gatctagcaa	180
tttaactctg	gagtccttca	agactgctgt	tcttacgaag	ttcacgtccc	tgaatcatgt	240
tctgtatg	aagcctgaaa	gacctcaa	tctaaaaggt	ggcgataaat	tgaagggtta	300
caaaatatac	cctgcgggct	tgacacagag	gcaagctctt	tataccttcc	agttcaacgg	360
ggatgttgat	ttcagaagtc	acttgagag	caatccttgt	gccaagtttg	aagtaatttt	420
tgtgtagcat	atgttgagct	acctacaatt	tacatgatca	cctagcatta	gctctttcac	480
ttaactgaga	gaatgaagtt	ttaggaatga	gtatgaccat	ggagtcggca	tggctttgta	540
atgcctaccc	tactttggcc	aactcatcgg	ggattttacat	tcagaaaata	tacatgactt	600
caaccatact	taaaccctt	tttgttaagat	aactgaatgt	tcataatttaa	tgttgggttg	660
tagtgttttt	acttgattat	atccagacag	ttacaagttg	gacaacaaga	ttgtgggtct	720
gtactgttat	ttattttatt	tttttttagc	agaaacacct	tatcttttgt	ttcgtttgaa	780
tgtagaatga	aaataaaaaga	aagaaaatat	aacatcatcg	gccgcgcttg	tctaatttcg	840
ggcagttagg	atcctctccg	gtcaccggaa	agtttcagta	gaagaaacaa	aacaccgtga	900
ctaaaatgat	actattattt	tatttattgt	gtttttcttt	tttctaccgg	aacttttttag	960
aacggatccc	aactcgttcc	ggggccgcta	caactgaaac	aaaagaagat	attttctctc	1020
tcttcagaaa	tgtaagtttt	cctttacaga	taccattca	ccatttgatt	catagtgtgt	1080
gactagagat	aaagcatact	aatttgactc	ttggaaaccc	ataaagttaa	tgttatccgt	1140
gttctggacc	aatccacttg	ggggcataac	ctgtgtctat	gtgtggtttg	gtttccattc	1200
tgatttatgc	ggcgacttgt	aattttaa	ctaggagggg	cagacattga	acaatcccaa	1260
tattttaata	acttatgcaa	gatttttttt	attaatgaga	tgatgtgttt	gtgactgaga	1320
ttgagtcata	catttcacta	agaaatggtt	ccaagtacca	aactatcatg	accagttgc	1380
aaacatgacg	ttcgggagtg	gtcactttga	tagttcaatt	tcactctggc	ttcttattcc	1440
ttttataatt	ctaattcttc	ttgtgtaaac	tatttcatgt	attatttttc	tttaaaattt	1500
acatgtcatt	tattttgcct	cactaactca	attttgcata	taacaatgat	aagtgatatt	1560
ttgactcaca	aaattttacat	caaattttcga	catcgtttat	tatgttcatt	ggatgattaa	1620
caaatataac	aaactttgca	actaatatac	caccaactga	atataattaa	ctataactgt	1680
gaaagtagtt	aaccatattt	tttagatgta	tatatcatcc	gttgaatgta	attattcata	1740
tatttgaact	aagttaccct	acaacttaaa	gaacttaag	aactcggttt	gagacctggg	1800
gacgaaaatg	taatgagact	ttaatgttga	ctttgacacc	gcaccacatg	tgcccttttac	1860
atatagttaa	tatgacaagt	aatgacaatc	cttgccttat	tataaggcga	cccttagctc	1920
caaccaaagg	acgatggagt	taagaaagaa	actcttgctt	acttgaagg	tccacacttc	1980
ttcactcacc	tctcaatttc	atcctacaaa	aatgtccaaa	cttctctttc	tcacaatcac	2040
aaactcattc	caaacacact	ctcttctcca	aaaatgtctt	ccttctcagt	acgtttcctc	2100
acccaccat	ccatctctcg	tcccaacaaa	acatggctac	tatctgctgc	aactccatca	2160
gttgc						2165

<210> 10

<211> 1316

<212> DNA

<213> Artificial Sequence

<220>

<223> Sequences to be used as promoter for regulating
expression of foreign genes

<400> 10

gatcctctcc	ggtcaccgga	aagtttcagt	agaagaaaca	aaacaccgtg	actaaaatga	60
tactattatt	ttatttattg	tgtttttctt	ttttctaccg	gaacttttta	gaacggatcc	120
caactcgttc	cggggccgct	acaactgaaa	caaaagaaga	tattttctct	ctcttcagaa	180
atgtaagttt	tccttttacag	ataccatttc	accatttgat	tcagatgtgg	tgactagaga	240
taaagcatac	taatttgact	cttggaacc	cataaagttt	atggtatccg	tgttctggac	300
caatccactt	gggggcataa	cctgtgtcta	tgtgtggttt	ggtttccatt	ctgatttatg	360
cggcgacttg	taatttaaaa	tctaggaggg	gcagacattg	aacaatccca	atattttaat	420
aacttatgca	agattttttt	tattaatgag	atgatgtgtt	tgtgactgag	attgagtcac	480
acatttcact	aagaaatggg	tccaagtacc	aaactatcat	gaccagttg	caaacatgac	540
gttcgggagt	ggtcactttg	atagttcaat	ttcatcttgg	cttcttattc	cttttataat	600
tctaattcct	cttggtgtaa	ctatttcgat	tattattttt	ctttaaaatt	tacatgtcat	660
ttattttgcc	tcactaactc	aattttgcat	ataacaatga	taagtgatat	tttgactcac	720
aaaattttaca	tcaaatttcg	acatcgttta	ttatgttcat	tggatgatta	acaaatataa	780
caaactttgc	aactaattaa	ccaccaactg	aatataatta	actataactg	tgaaagtagt	840
taaccatatt	ttttagatgt	atatatcatc	cgttgaatgt	aattattcat	atatttgaac	900
taagttaccc	tacaacttaa	agaacttaaa	gaactcgggt	tgagacctgg	ggacgaaaat	960
gtaatgagac	tttaatgttg	actttgacac	cgcaccacat	gtgcctttta	catatagttt	1020
atatgacaag	taatgacaat	ccttgctcta	ttataaggcg	acccttagct	ccaaccaaag	1080
gacgatggag	ttaagaaaga	aactcttgct	tacttgtaag	gtccacactt	cttcactcac	1140
ctctcaattt	catctacaa	aaatgtccaa	acttctcttt	ctcacaatca	caaactcatt	1200
ccaaacacac	tctcttctcc	aaaaatgtct	tccttctcag	tacgtttcct	cacccaccca	1260
tccatctctc	gtcccaacaa	aacatggcta	ctatctgctg	caactccatc	agttgc	1316

<210> 11

<211> 2971

<212> DNA

<213> Artificial Sequence

<220>

<223> Sequences to be used as promoter for regulating expression of foreign genes

<400> 11

ctgtacattc	atcttgccgc	ctttgcattc	acttggccac	aaagagtaga	gagaaggaag	60
agaagagccc	agacttcaag	aagcgacctt	gcaagtgcac	tcgagggtca	gaaactgtat	120
atcatatcta	tgtgagagaa	aggggaacat	tgagatgga	gtccatttac	ttgaggatata	180
cttattattt	tgatcaataa	atttgtatac	ttcttattta	gatcaataaa	tttgtcatta	240
agctataatc	caaaataaat	tacgatcaaa	tatgcaaagt	ttagccagta	cttgtgttaa	300
acttgatggc	atctcttggt	ttctttggca	atcacatgcc	taagaaataa	atagtatcat	360
atgatttgtt	ttggtcagac	ttcagagtca	gatgactctg	tttgataaa	cagcttaatt	420
aagcgcttat	agaatatcat	atgatttgtt	ttggtcagac	ttcagagcat	ctcttggttt	480
ctctggcaat	catatgccta	agaaataaat	agtatcatat	gatttgtgtt	ggtcagactt	540
cagagtcaga	tgaccctggt	tgggtaaaca	gcttaattaa	gtgcttatag	aataagcgct	600
tatcatataa	gtgcttttgt	acagttattt	ctatgaaagt	agaagaaata	gtcatattgt	660
tttaataata	gctatcctgg	agagcttggt	gaaataacca	gaaaagaact	tatggacacg	720
tcatgagctg	tttacataag	atctccctaa	cagtcctcaa	agtgtttatg	ccagtagata	780
aattcaaata	agtcaatcta	aacagaccct	aaatccatta	tggtagctat	cattttagct	840
tattccatct	ttattaagaa	tgtcatgaga	taaacacatt	attttgacac		900
aaatgggcag	atctagcaat	ttactctgg	agtccttcaa	gactgctgtt	cttacgaagt	960
tcacgtccct	gaatcatggt	cctgtatgga	agcctgaaag	acctcaaatt	ctaaaagggtg	1020
gcgataaatt	gaagggtttac	aaaatatacc	ctgcgggctt	gacacagagg	caagctcttt	1080
ataccttcca	gttcaacggg	gatgttgatt	tcagaagtca	cttgagagc	aatccttggtg	1140
ccaagtttga	agtaattttt	gtgtagcata	tgttgagcta	cctacaattt	acatgatcac	1200
ctagcattag	ctctttcact	taactgagag	aatgaagttt	taggaatgag	tatgaccatg	1260
gagtcggcat	ggctttgtaa	tgcctaccct	actttggcca	actcatcggg	gatttacatt	1320
cagaaaatat	acatgacttc	aaccataactt	aaaccccttt	ttgtaagata	actgaatggt	1380
catattttaat	gttgggttgt	agtgttttta	cttgattata	tccagacagt	tacaagttgg	1440
acaacaagat	tgtgggtctg	tactgttatt	tatttatttt	tttttttagca	gaaacacctt	1500

10/17

atcttttgtt	tcgtttgaat	gtagaatgaa	aataaaaagaa	agaaaatata	acatcatcgg	1560
ccgcgcttgt	ctaatttcgg	gcagtttaga	tcctctccgg	tcaccggaaa	gtttcagtag	1620
aagaaacaaa	acaccgtgac	taaaatgata	ctattatttt	atztatgtg	ttttctttt	1680
ttctaccgga	actttttaga	acggatccca	actcgttccg	gggccgctac	aactgaaaca	1740
aaagaagata	ttttctctct	cttcagaaat	gtaagttttc	ctttacagat	acccattcac	1800
catttgattc	agatgtggtg	actagagata	aagcatacta	atttgactct	tggaaacca	1860
taaagtttat	gttatccgtg	ttctggacca	atccacttgg	gggcataacc	tgtgtctatg	1920
tgtggtttg	tttccattct	gatttatgcg	gcgacttgta	atttaaaatc	taggaggggc	1980
agacattgaa	caatcccaat	attttaataa	cttatgcaag	atttttttta	ttaatgagat	2040
gatgtgtttg	tgactgagat	tgagtcatac	atttcactaa	gaaatgggtc	caagtaccaa	2100
actatcatga	cccagttgca	aacatgacgt	tcgggagtg	tcactttgat	agttcaattt	2160
catcttggct	tcttattcct	tttataattc	taattcttct	tgtgtaaact	atttcatgta	2220
ttatttttct	ttaaaattta	catgtcattt	attttgcttc	actaactcaa	ttttgcatat	2280
aacaatgata	agtgatattt	tgactcacaa	aattttacatc	aaatttcgac	atcgtttatt	2340
atgttcattg	gatgattaac	aaatataaca	aactttgcaa	ctaattaacc	accaactgaa	2400
tataattaac	tataactgtg	aaagtagtta	accatatttt	ttagatgtat	atatcatccg	2460
ttgaatgtaa	ttattcatat	atttgaacta	agttacccta	caacttaaag	aacttaaaga	2520
actcggtttg	agacctgggg	acgaaaatgt	aatgagactt	taatgttgac	tttgacaccg	2580
caccacatgt	gccttttaca	tatagtttat	atgacaagta	atgacaatcc	ttgctctatt	2640
ataaggcgac	ccttagctcc	aaccaaagga	cgatggagtt	aagaaagaaa	ctcttgctta	2700
cttgtaaggt	ccacacttct	tcactcacct	ctcaatttca	tcctacaaaa	atgtccaaac	2760
ttctctttct	cacaatcaca	aactcattcc	aaacacactc	tcttctccaa	aaatgtcttc	2820
cttctcagta	cgtttctcca	ccccaccatc	catctctcgt	cccaacaaaa	catggctact	2880
atctgctgca	actccatcag	ttgcacctgt	ttcaacacca	caagttgatg	catcaaggtt	2940
ggagcctaga	gttgaggaaa	aagatggtta	c			2971

<210> 12

<211> 2232

<212> DNA

<213> Artificial Sequence

<220>

<223> Sequences to be used as promoter for regulating
expression of foreign genes

<400> 12

gatctcccta	acagtctcaa	aagtgtttat	gccagtagat	aaattcaa	aagtcaatct	60
aaacagaccc	taaatccatt	atggtacctt	tcatttttagc	ttattccatc	tttattaaga	120
atgtcatgag	ataacataat	gataacacat	tattttgaca	caaatgggca	gatctagcaa	180
tttaactctg	gagtccttca	agactgctgt	tcttacgaag	ttcacgtccc	tgaatcatgt	240
tcctgtatgg	aagcctgaaa	gacctcaa	tctaaaagg	ggcgataaat	tgaagggtta	300
caaaatatac	cctgcgggct	tgacacagag	gcaagctctt	tataccttcc	agttcaacgg	360
ggatgttgat	ttcagaagtc	acttgagag	caatccttgt	gccaagtttg	aagtaatttt	420
tgtgtagcat	atgttgagct	acctacaatt	tacatgatca	cctagcatta	gctctttcac	480
ttaaactgaga	gaatgaagtt	ttaggaatga	gtatgaccat	ggagtcggca	tggctttgta	540
atgcctaccc	tactttggcc	aactcatcgg	ggatttacat	tcagaaaata	tacatgactt	600
caaccatact	taaacccctt	tttgtaagat	aactgaatgt	tcataattta	tgttgggttg	660
tagtggtttt	acttgattat	atccagacag	ttacaagttg	gacaacaaga	ttgtgggtct	720
gtactgttat	ttattttatt	tttttttagc	agaaacacct	tatcttttgt	ttcgtttgaa	780
tgtagaatga	aaataaaaaga	aagaaaatat	aacatcatcg	gccgcgcttg	tctaatttcg	840
ggcagttagg	atcctctccg	gtcaccggaa	agtttcagta	gaagaaacaa	aacaccgtga	900
ctaaaatgat	actattat	tattttattg	gtttttcttt	tttctaccgg	aacttttttag	960
aacggatccc	aactcgttcc	ggggccgcta	caactgaaac	aaaagaagat	attttctctc	1020
tcttcagaaa	tgtaagtttt	cctttacaga	taccatttca	ccatttgatt	cagatgtggg	1080
gactagagat	aaagcatact	aatttgactc	ttggaaaccc	ataaagttta	tgttatccgt	1140
gttctggacc	aatccacttg	ggggcataac	ctgtgtctat	gtgtgggttg	gtttccattc	1200
tgatttatgc	ggcgacttgg	aatttaaaat	ctaggagggg	cagacattga	acaatcccaa	1260
tatttttaata	acttatgcaa	gatttttttt	attaatgaga	tgatgtgttt	tgactgaga	1320
ttgagtcata	catttcacta	agaaatgggt	ccaagtacca	aactatcatg	acccagttgc	1380
aaacatgacg	ttcgggagtg	gtcactttga	tagttcaatt	tcactctggc	ttcttattcc	1440

ttttataatt	ctaattcttc	ttgtgtaa	ac	tatttcatgt	attatttttc	tttaaaattt	1500
acatgtcatt	tattttgcct	cactaactca	at	ttttgcata	taacaatgat	aagtgatatt	1560
ttgactcaca	aaattttacat	caaatttcga	cat	cgttttat	tatgttcatt	ggatgattaa	1620
caaataataac	aaacttttgca	actaattaac	caccaactga	atataattaa	ctataactgt		1680
gaaagtagtt	aaccatattt	tttagatgta	tatatcatcc	gttgaatgta	attattcata		1740
tatttgaact	aagttaccct	acaacttaaa	gaacttaaa	gaactcggttt	gagacctggg		1800
gacgaaaaatg	taatgagact	ttaatgttga	ctttgacacc	gcaccacatg	tgcccttttac		1860
atatagttta	tatgacaagt	aatgacaatc	cttgctctat	tataaggcga	cccttagctc		1920
caaccaaagg	acgatggagt	taagaaagaa	actcttgctt	acttgtaagg	tccacacttc		1980
ttcactcacc	tctcaatttc	atcctacaaa	aatgtccaaa	cttctctttc	tcacaatcac		2040
aaactcattc	caaacacact	ctcttctcca	aaaatgtctt	ccttctcagt	acgtttcctc		2100
acccaccat	ccatctctcg	tcccaacaaa	acatggctac	tatctgctgc	aactccatca		2160
gttgacactg	tttcaacacc	acaagttgat	gcatcaagg	tggagcctag	agttgaggaa		2220
aaagatgggt	ac						2232

<210> 13

<211> 1383

<212> DNA

<213> Artificial Sequence

<220>

<223> Sequences to be used as promoter for regulating expression of foreign genes

<400> 13

gacacctctc	ggtcaccgga	aagtttcagt	agaagaaaca	aaacaccgtg	actaaaatga	60
tactattatt	ttattttattg	tgtttttctt	ttttctaccg	gaacttttta	gaacggatcc	120
caactcgttc	cggggccgct	acaactgaaa	caaaagaaga	tattttctct	ctcttcagaa	180
atgtaagttt	tcctttacag	ataccattc	accatttgat	tcagatgtgg	tgactagaga	240
taaagcatac	taatttgact	cttggaacc	cataaagttt	atgttatccg	tggtctggac	300
caatccactt	gggggcataa	cctgtgtcta	tgtgtggttt	ggtttccatt	ctgatttatg	360
cggcgacttg	taatttaaaa	tctaggaggg	gcagacattg	aacaatccca	atattttaat	420
aacttatgca	agattttttt	tattaatgag	atgatgtgtt	tgtgactgag	attgagtcac	480
acattttcact	aagaaatggg	tccaagtacc	aaactatcat	gacccagttg	caaacatgac	540
gttcgggagt	ggtcactttg	atagttcaat	ttcatcttgg	cttcttattc	cttttataat	600
tctaattctt	cttgtgtaaa	ctattttcatg	tattattttt	ctttaaaatt	tacatgtcat	660
ttattttgcc	tactaaactc	aattttgcat	ataacaatga	taagtgatat	tttgactcac	720
aaaattttaca	tcaaatttcg	acatcgttta	ttatgttcat	tggatgatta	acaaatataa	780
caaactttgc	aactaattaa	ccaccaactg	aatataatta	actataactg	tgaaagtagt	840
taaccatatt	tttttagatgt	atatatcatc	cgttgaatgt	aattattcat	atatttgaac	900
taagttaccc	tacaacttaa	agaacttaaa	gaactcgggt	tgagacctgg	ggacgaaaat	960
gtaatgagac	tttaatgttg	actttgacac	cgcaccacat	gtgcctttta	catatagttt	1020
atatgacaag	taatgacaat	ccttgctcta	ttataaggcg	acccttagct	ccaaccaaag	1080
gacgatggag	ttaagaaaga	aactcttgct	tacttgtaag	gtccacactt	cttcaactcac	1140
ctctcaattt	catcctacaa	aaatgtccaa	acttctcttt	ctcacaatca	caaactcatt	1200
ccaaacacac	tctcttctcc	aaaaatgtct	tccttctcag	tacgtttcct	caccccacca	1260
tccatctctc	gtcccaacaa	aacatggcta	ctatctgctg	caactccatc	agttgcacct	1320
gtttcaacac	cacaagttga	tgcacaaagg	ttggagccta	gagttgagga	aaaagatggg	1380
tac						1383

<210> 14

<211> 3472

<212> DNA

<213> Artificial Sequence

<220>

<223> Sequences to be used as terminator for regulating expression of foreign genes

<400> 14

caagttgatg	catcaaggtt	ggagcctaga	gttgaggaaa	aagatgggta	ctgggttttg	60
aaggaagagt	atagaggagg	tattaatcct	caggagaaa	gaaagaacct	120	
atgaagcttt	ttatggaagg	tgggattaat	gatttggtca	atatgtctct	tgaagagatt	180
gaaagctcta	agcttactaa	agatgatatt	gatgttagac	ttaaatggct	tgggtctttt	240
catagaagga	aacatcattg	taagtttttt	tactttcttt	ttatacttca	aagtctctct	300
atactctgta	tttggtttatt	agttttttgt	gacttaaata	ttctctttga	tttacatagt	360
gaaactccat	tcttggtttc	gaaattgtag	tgtgtatagt	ctagaaaatt	aagaagtaga	420
caaaatgatt	tatgagattg	taaattgtag	gctttttatc	aattttattaa	tttagagac	480
caaaatttgc	ctatcttatt	tggaccaatt	tattgtatgt	taggatcgac	atgagtttag	540
caaaatcatg	acggcaccat	gactgtgttg	aagcttcttt	gtgtaacttt	aaccaaaatt	600
atatggcaca	ccatgattat	gcaaactcac	cgtcaatcca	aacatagaaa	ttcagtggtta	660
atctttgtga	caataaaaaa	ctatgagtta	tgttgtagta	atcttattcc	attgtgaaac	720
tcagatggta	gatttatgat	gagactaaaa	ctcccaaatg	gggtaacaac	aagtgtctca	780
acaagatact	tggcgagtgt	gataaaaaaa	tatggcaaag	acggatgtgc	tgatgtgaca	840
acgaggcaga	attggcaaat	tcgaggtgta	acgttacctg	atgtccctga	aattcttaag	900
ggccttgcag	aggctcggtt	gacaagtctg	cagagtggaa	tggacaatgt	tcgaaaccca	960
gttggttaacc	ctcttgctgg	tattgacctt	gatgagattg	ttgatacaag	accttacacc	1020
aatttgctgt	cccaattcat	caactgcta	tcacttggtg	atccaaccat	tacaaacttg	1080
taagtctaaa	ctatctcatc	tttatatttc	actcattata	tcataattag	agttagttac	1140
ttgcattgca	agcattacgt	gaccgtgtgt	agcctctaaa	tccttttgat	aatatgtgca	1200
ggccaaggaa	gtggaatgta	tgtgtgatag	gttcccatga	tcttttcgag	catccgcata	1260
ttaacgatct	tgccttatatg	cctgctaata	aggatggctg	atgttgattc	aacttattgg	1320
tgggtgggtt	ctttagtccc	aagcgatgtg	ctgaagcagt	tcacttgat	gcattgggtct	1380
ctgcagatga	tgttatccca	ctttgtaaag	ctgtccttga	gacctatagg	gacctcggca	1440
caagagggaa	tagacagaaa	accagaatga	tgtggttgat	cgatgaactt	gtaagtacc	1500
actttttttc	ttcacatatt	attaactgaa	gtgactttta	cgaccatttt	acaattgaaa	1560
tttaagtga	ttttagccct	atcattacaa	gaacaaattt	gttaattcac	tagcaagagc	1620
aattccactt	tggcttggac	atgacaagtg	tttgtgaaat	gcaggggata	gaagtattca	1680
gatcagaggt	ggaaaaaaga	atgccagaga	agaagctaga	gagagcatcc	aaagaagaac	1740
ttgtccaaaa	acaatggaaa	gaggagacat	cttaggtgtt	catccacaaa	aacaagaagg	1800
tttaagctat	gttggaaattc	acattccagt	tggtagaatc	caagcagatg	agatggaaga	1860
gctagctcgt	atcgccgatg	aatacggaa	cggagaacta	aggctaaccg	tggagcaaaa	1920
cataataatt	ccaaatgtgg	aaaactcaaa	acttgatgca	ttgctaaatg	aacctctctt	1980
gaaagacaaa	ttctcaccag	aaccttccat	cctaataaaa	acacttggtg	catgcactgg	2040
taaccaattt	tgtggccaag	caataattga	aacaaaacaa	agagctttta	aagtaactga	2100
agaagttgaa	agacatgtgg	ctgtgagcaa	accagtgaga	atgcattgga	ctggttgtcc	2160
taacacttgt	ggtcaagttc	aggttgctga	tattgggttt	atgggttgta	tggctaggga	2220
tgagaatgg	aaggctactg	aaggtgttga	tattttcctt	ggtgggagaa	ttggaagtga	2280
ttctcattta	gctgaggtgt	ataagaaaag	tgtcccttgc	aaggacttgg	tgcctattgt	2340
agctgatatt	ttgggttaaat	attttggagc	tgtccaaaag	aatagagaag	aaggggatga	2400
ttaaagtata	taggtatttg	gtgattttta	ttgctcttac	acaaaattat	tatgttctgt	2460
ccaaaatata	aagtcacaag	ggataattga	gattgagatg	cagcacgcca	cacatgaact	2520
tgtacatttg	gataagtcac	ttttcattgc	tattttataa	gttacacttt	gaattttata	2580
ataaatttta	ttttatttca	aggaccagat	tttataagga	aaccgcta	ctaactatct	2640
ttactcgtaa	tttgtcattt	gagagctacg	gagatcgttg	agtttacgta	tgagtgttta	2700
gtctcacatt	aattatgaat	ggtcaaaatg	ttaaatttat	aagagatgta	atctatatac	2760
ctaattgcatt	aaaaatttgg	atggagatgc	gacgcccccc	ttttttgtgg	tcctgaagta	2820
tagacttggt	gtcgtctctg	gtgcactctc	atacttccca	acaaggagaa	aaaactacca	2880
taacaattaa	caaactaaca	tttggttattt	aaaaaaacat	acggatactg	ttttttcccc	2940
atattattagg	aagatgatgg	cttggttttc	aatggctgag	tttttttttt	tttttgctgg	3000
gagttgaagt	atcgggaaaa	ctaaatatgc	tatgacttta	aacatttgtgt	tgatatatga	3060
ttagttttca	acttacttaa	aaagtggcaa	actagtttag	tggttctctc	ccttctctgt	3120
agttcaagga	acatgggttt	gaactctgtc	caaatttttg	tactttcaat	tatccataca	3180
tttaaaagct	atataccaca	tcatttatatt	caagtcaatg	atcatgcggc	ctgccacatt	3240
agcatcgatg	tacacattaa	ttttaagtgg	catgaacaca	ttaacatttc	ataaaagcta	3300
tgtgccagat	catcattcaa	gtctatgcac	acatgggtcaa	cacatttagta	ccattttttt	3360
ttttatttgt	gatcagattg	gatgtcggta	ttgttgtgat	gctacaaact	caaacaatct	3420
ccagctgtta	gagaacgtcg	aaaatgaatg	catcacgggt	gcacacttag	at	3472

13/17

<211> 617
 <212> DNA
 <213> Artificial Sequence

<220>

<223> Sequences to be used as terminator for regulating
 expression of foreign genes

<400> 15

agtatatagg	tatttggtga	ttttaattgc	ctctacacaa	aattattatg	ttctgtccaa	60
aatataaagt	cacaagggat	aattgagatt	gagatgcagc	acgccacaca	tgaacttgta	120
catttggtgata	agtcattttt	cattgctatt	ttataagtta	cactttgaat	tttataataa	180
atttttatttt	atttcaagga	ccagatttta	taaggaaacc	gctaatactaa	ctatctttac	240
tcgtaatttg	tcatttgaga	gctacggaga	tcgttgagtt	tacgtatgag	tgtttagtct	300
cacattaatt	atgaatggtc	aaaatgttaa	atttataaga	gatgtaatct	atatacctaa	360
tgcatataaaa	atttggatgg	agatgcgacg	ccccctttt	ttgtgggtcct	gaagtataga	420
cttggtgtcg	cttctgggtgc	actctcatac	ttcccaacaa	ggagaaaaaa	ctaccataac	480
aattaacaaa	ctaacatttg	ttattttaaaa	aaacatacgg	atactgtttt	ttccccattt	540
attaggaaga	tgatggcttg	gatttcaatg	gctgagttta	tttttttttt	ggtcgggagt	600
tgaagtatcg	ggaaaaac					617

<210> 16
 <211> 502
 <212> DNA
 <213> Artificial Sequence

<220>

<223> Sequences to be used as terminator for regulating
 expression of foreign genes

<400> 16

ttgtacattt	ggataagtca	tttttcattg	ctattttata	agttacactt	tgaattttat	60
aataaatttt	attttatttc	aaggaccaga	ttttataagg	aaaccgctaa	tctaactatc	120
tttactcgta	atttgtcatt	tgagagctac	ggagatcggt	gagtttacgt	atgagtgttt	180
agtctcacat	taattatgaa	tggtcaaaat	gttaaattta	taagagatgt	aatctatata	240
cctaattgcat	taaaaatttg	gatggagatg	cgacgcccc	cttttttggtg	gtcctgaagt	300
atagacttgt	tgctgcttct	ggtgcactct	catacttccc	aacaaggaga	aaaaactacc	360
ataacaatta	acaaactaac	atttgttatt	taaaaaaaaca	tacggatact	gttttttccc	420
catttattag	gaagatgatg	gcttggattt	caatggctga	gtttattttt	tttttggtcg	480
ggagttgaag	tatcgggaaa	ac				502

<210> 17
 <211> 25
 <212> DNA
 <213> Artificial Sequence

<220>

<223> Sequences to be used as primers

<400> 17

gatattgatg ttgactcaa gtggc

25

<210> 18
 <211> 21
 <212> DNA
 <213> Artificial Sequence

<220>

<223> Sequences to be used as primers

<400> 18

cacysattcc acttcctwgg c

21

<210> 19

<211> 27

<212> DNA

<213> Artificial Sequence

<220>

<223> Sequences to be used as primers

<400> 19

ttgtcacatc agcacatccg tctttgc

27

<210> 20

<211> 27

<212> DNA

<213> Artificial Sequence

<220>

<223> Sequences to be used as primers

<400> 20

tcgccaagta tcttgtttga gcacttg

27

<210> 21

<211> 27

<212> DNA

<213> Artificial Sequence

<220>

<223> Sequences to be used as primers

<400> 21

agagcccggg agaagagagt gtgtttg

27

<210> 22

<211> 28

<212> DNA

<213> Artificial Sequence

<220>

<223> Sequences to be used as primers

<400> 22

ttctcccggg ggacgagaga tggatggt

28

<210> 23
<211> 30
<212> DNA
<213> Artificial Sequence

<220>
<223> Sequences to be used as primers

<400> 23
ttctcccggg gttgaaacag gtgcaactga 30

<210> 24
<211> 28
<212> DNA
<213> Artificial Sequence

<220>
<223> Sequences to be used as primers

<400> 24
ttctcccggg taaccatctt tttcctca 28

<210> 25
<211> 24
<212> DNA
<213> Artificial Sequence

<220>
<223> Sequences to be used as primers

<400> 25
cacacttctt cactcacctc tcaa 24

<210> 26
<211> 21
<212> DNA
<213> Artificial Sequence

<220>
<223> Sequences to be used as primers

<400> 26
atctaggagg ggcagacatt g 21

<210> 27
<211> 24
<212> DNA
<213> Artificial Sequence

<220>
<223> sequences to be used as primers

<400> 27

tcggtataaa gacttcgcgc tgat

24

<210> 28

<211> 27

<212> DNA

<213> Artificial Sequence

<220>

<223> sequences to be used as primers

<400> 28

atgtcttcct tctcagtagc tttcctc

27

<210> 29

<211> 30

<212> DNA

<213> Artificial Sequence

<220>

<223> sequences to be used as primers

<400> 29

caagttgatg catcaaggtg ggagcctaga

30

<210> 30

<211> 30

<212> DNA

<213> Artificial Sequence

<220>

<223> sequences to be used as primers

<400> 30

agaagagctc agtatatagg tatttggtga

30

<210> 31

<211> 30

<212> DNA

<213> Artificial Sequence

<220>

<223> sequences to be used as primers

<400> 31

agaagagctc ttgtacattt ggataagtca

30

<210> 32

<211> 30

<212> DNA

<213> Artificial Sequence

<220>

<223> sequences to be used as primers

<400> 32

agaagaattc gttttcccga tacttcaact

30